WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: C12N 15/82, 15/79, 15/54, A01H 5/00 (11) International Publication Number:

WO 97/04113

(43) International Publication Date:

6 February 1997 (06.02.97)

(21) International Application Number:

PCT/EP96/03053

A2

(22) International Filing Date:

12 July 1996 (12.07.96)

(30) Priority Data:

9514437.4

14 July 1995 (14.07.95)

GB

(71) Applicant (for all designated States except US): DANISCO A/S [DK/DK]; Langebrogade 1, P.O. Box 17, DK-1001 Copenhagen K (DK).

(72) Inventor; and

(75) Inventor/Applicant (for US only): POULSEN, Peter [DK/DK]; Holsteinsgade 52, 3, DK-2100 Copenhagen K (DK).

(74) Agents: HARDING, Charles, Thomas et al.; D. Young & Co., 21 New Fetter Lane, London EC4A 1DA (GB).

(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY. CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published

Without international search report republished upon receipt of that report.

With an indication in relation to a deposited microorganism furnished under Rule 13bis separately from the description.

Date of receipt by the International Bureau:

20 August 1996 (20.08.96)

(54) Title: INHIBITION OF GENE EXPRESSION

(57) Abstract

A method of inhibiting gene expression is described. The method, which affects enzymatic activity in a plant, comprises expressing in a plant (or a cell, a tissue or an organ thereof) a nucleotide sequence wherein the nucleotide sequence codes, partially or completely, for an intron in a sense orientation; and wherein the nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
ΑŲ	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	· IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgystan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic	SD	Sudan
CF	Central African Republic		of Korea	SE	Sweden
CG	Congo	· KR	Republic of Korea	SG	Singapore
CH	Switzerland	· KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LR	Liberia	SZ	Swaziland
CS	Czechoslovakia	LT	Lithuania .	TD	Chad
CZ	Czech Republic	LU	Luxembourg	TG	Togo
DE	Germany	LV	Latvia	TJ	Tajikistan
DK	Denmark	MC	Monaco	TT	Trinidad and Tobago
EE	Estonia	MD	Republic of Moldova	UA	Ukraine
ES .	Spain	MG	Madagascar	UG	Uganda
FI.	Finland	ML	Mali	US	United States of America
FR	Prance	MN	Mongolia	UZ	Uzbekistan
GA	Gabon	MR	Mauritania	VN	Viet Nam

10

- 15

20

25

30

INHIBITION OF GENE EXPRESSION

The present invention relates to a method of inhibiting gene expression, particularly inhibiting gene expression in a plant. The present invention also relates to a nucleotide sequence useful in the method. In addition, the present invention relates to a promoter that is useful for expressing the nucleotide sequence.

Starch is one of the main storage carbohydrates in plants, especially higher plants. The structure of starch consists of amylose and amylopectin. Amylose consists essentially of straight chains of α -1-4-linked glycosyl residues. Amylopectin comprises chains of α -1-4-linked glycosyl residues with some α -1-6 branches. The branched nature of amylopectin is accomplished by the action of *inter alia* an enzyme commonly known as the starch branching enzyme ("SBE"). SBE catalyses the formation of branch points in the amylopectin molecule by adding α -1,4 glucans through α -1,6-glucosidic branching linkages. The biosynthesis of amylose and amylopectin is schematically shown in Figure 1, whereas the α -1-4-links and the α -1-6 links are shown in Figure 2.

It is known that starch is an important raw material. Starch is widely used in the food, paper, and chemical industries. However, a large fraction of the starches used in these industrial applications are post-harvest modified by chemical, physical or enzymatic methods in order to obtain starches with certain required functional properties.

Within the past few years it has become desirable to make genetically modified plants which could be capable of producing modified starches which could be the same as the post-harvest modified starches. It is also known that it may be possible to prepare such genetically modified plants by expression of antisense nucleotide coding sequences. In this regard, June Bourque provides a detailed summary of antisense strategies for the genetic manipulations in plants (Bourque 1995 Plant Science 105 pp 125-149).

10

15

20

25

30

Whilst it is known that enzymatic activity can be affected by expression of particular nucleotide sequences (for example see the teachings of Finnegan and McElroy [1994] Biotechnology 12 883-888; and Matzke and Matzke [1995] TIG 11 1-3) there is still a need for a method that can more reliably and/or more efficiently and/or more specifically affect enzymatic activity.

According to a first aspect of the present invention there is provided a method of affecting enzymatic activity in a plant (or a cell, a tissue or an organ thereof) comprising expressing in the plant (or a cell, a tissue or an organ thereof) a nucleotide sequence wherein the nucleotide sequence partially or completely codes (is) an intron in a sense orientation; and wherein the nucleotide sequence does not contain a sequence that is a sense exon sequence normally associated with the intron.

According to a second aspect of the present invention there is provided a method of affecting enzymatic activity in a starch producing organism (or a cell, a tissue or an organ thereof) comprising expressing in the starch producing organism (or a cell, a tissue or an organ thereof) a nucleotide sequence wherein the nucleotide sequence codes, partially or completely, for an intron in a sense orientation; wherein the nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron; and wherein starch branching enzyme activity is affected and/or the levels of amylopectin are affected and/or the composition of starch is changed.

According to a third aspect of the present invention there is provided a sequence comprising the nucleotide sequence shown as any one of SEQ.I.D. No. 1 to SEQ.I.D. No. 13 or a variant, derivative or homologue thereof.

According to a fourth aspect of the present invention there is provided a promoter comprising the sequence shown as SEQ.I.D. No. 14 or a variant, derivative or homologue thereof.

25

According to a fifth aspect of the present invention there is provided a construct capable of comprising or expressing the present invention.

According to a sixth aspect of the present invention there is provided a vector comprising or expressing the present invention.

According to a seventh aspect of the present invention there is provided a cell, tissue or organ comprising or expressing the present invention.

- According to an eighth aspect of the present invention there is provided a transgenic starch producing organism comprising or expressing the present invention. According to a ninth aspect of the present invention there is provided a starch obtained from the present invention.
- According to a tenth aspect of the present invention there is provided pBEA11 (NCIMB 40754). According to an eleventh aspect of the present invention there is provided a sense nucleotide sequence that is obtainable from λ -SBE 3.2 (NCIMB 40751) or λ -SBE 3.4 (NCIMB 40752) or a variant, derivative or homologue thereof.
- A key advantage of the present invention is that it provides a method for preparing modified starches that is not dependent on the need for post-harvest modification of starches. Thus the method of the present invention obviates the need for the use of hazardous chemicals that are normally used in the post-harvest modification of starches.

In addition, the present invention provides *inter alia* genetically modified plants which are capable of producing modified and/or novel and/or improved starches whose properties would satisfy various industrial requirements.

Thus, the present invention provides a method of preparing tailor-made starches in plants which could replace the post-harvest modified starches.

PCT/EP96/03053

WO 97/04113

Also, the present invention provides a method that enables modified starches to be prepared by a method that can have a more beneficial effect on the environment than the known post-harvest modification methods which are dependent on the use of hazardous chemicals and large quantities of energy.

5

10

15

An other key advantage of the present invention is that it provides a method that may more reliably and/or more efficiently and/or more specifically affect enzymatic activity when compared to the known methods of affecting enzymatic activity. With regard to this advantage of the present invention it is to be noted that there is some degree of homology between coding regions of SBEs. However, there is little or no homology with the intron sequences of SBEs. Thus, sense intron expression provides a mechanism to affect selectively the expression of a particular SBE. This advantageous aspect could be used, for example, to reduce or eliminate a particular SBE enzyme and replace that enzyme with another enzyme which can be another branching enzyme or even a recombinant version of the affected enzyme or even a hybrid enzyme which could for example comprise part of a SBE enzyme from one source and at least a part of another SBE enzyme from another source. This particular feature of the present invention is covered by the combination aspect of the present invention which is discussed in more detail later.

20

Thus the present invention provides a mechanism for selectively affecting SBE activity. This is in contrast to the prior art methods which are dependent on the use of for example antisense exon expression whereby it would not be possible to introduce new SBE activity without affecting that activity as well.

25

Preferably with the first aspect of the present invention starch branching enzyme activity is affected and/or wherein the levels of amylopectin are affected and/or the composition of starch is changed.

30

Preferably with the first or second aspect of the present invention the nucleotide sequence does not contain a sequence that is sense to an exon sequence.

Preferably with the fourth aspect of the present invention the promoter is in combination with a gene of interest ("GOI").

Preferably the enzymatic activity is reduced or eliminated.

5

20

Preferably the nucleotide sequence codes for at least substantially all of at least one intron in a sense orientation.

Preferably the nucleotide sequence codes, partially or completely, for two or more introns and wherein each intron is in a sense orientation.

Preferably the nucleotide sequence comprises at least 350 nucleotides (e.g. 350 bp), more preferably at least 500 nucleotides (e.g. 500 bp).

Preferably the nucleotide sequence comprises the sequence shown as any one of SEQ. I.D. No. 1 to SEQ.I.D. No. 13 or a variant, derivative or homologue thereof, including combinations thereof.

Preferably the nucleotide sequence is expressed by a promoter having a sequence shown as SEQ. I.D. No. 14 or a variant, derivative or homologue thereof.

A preferred aspect of the present invention therefore relates to a method of affecting

Preferably the transgenic starch producing organism is a plant.

enzymatic activity in a plant (or a cell, a tissue or an organ thereof) comprising expressing in the plant (or a cell, a tissue or an organ thereof) a nucleotide sequence wherein the nucleotide sequence codes, partially or completely, for an intron in a sense orientation; wherein the nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron; and wherein starch branching enzyme activity is affected and/or the levels of amylopectin are affected and/or the composition of starch is changed.

A more preferred aspect of the present invention therefore relates to a method of affecting enzymatic activity in a plant (or a cell, a tissue or an organ thereof) comprising expressing in the plant (or a cell, a tissue or an organ thereof) a nucleotide sequence wherein the nucleotide sequence codes, partially or completely, for an intron in a sense orientation; wherein the nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron; wherein starch branching enzyme activity is affected and/or the levels of amylopectin are affected and/or the composition of starch is changed; and wherein the nucleotide sequence comprises the sequence shown as any one of SEQ.I.D. No. 1 to SEQ.I.D. No. 13 or a variant, derivative or homologue thereof, including combinations thereof.

The term "nucleotide" in relation to the present invention includes DNA and RNA. Preferably it means DNA, more preferably DNA prepared by use of recombinant DNA techniques.

15

20

25

30

10

5

The term "intron" is used in its normal sense as meaning a segment of nucleotides, usually DNA, that does not encode part or all of an expressed protein or enzyme.

The term "exon" is used in its normal sense as meaning a segment of nucleotides, usually DNA, encoding part or all of an expressed protein or enzyme.

Thus, the term "intron" refers to gene regions that are transcribed into RNA molecules, but which are spliced out of the RNA before the RNA is translated into a protein. In contrast, the term "exon" refers to gene regions that are transcribed into RNA and subsequently translated into proteins.

The terms "variant" or "homologue" or "fragment" in relation to the nucleotide sequence of the present invention include any substitution of, variation of, modification of, replacement of, deletion of or addition of one (or more) nucleic acid from or to the respective nucleotide sequence providing the resultant nucleotide sequence can affect enzyme activity in a plant, or cell or tissue thereof, preferably wherein the resultant nucleotide sequence has at least the same effect as any one of

15

20

25

30

the sense sequences shown as SEQ.I.D. No.s 1-13. In particular, the term "homologue" covers homology with respect to similarity of structure and/or similarity of function providing the resultant nucleotide sequence has the ability to affect enzymatic activity in accordance with the present invention. With respect to sequence homology (i.e. similarity), preferably there is more than 80% homology, more preferably at least 85% homology, more preferably at least 90% homology, even more preferably at least 95% homology, more preferably at least 98% homology. The above terms are also synonymous with allelic variations of the sequences.

Likewise, the terms "variant" or "homologue" or "fragment" in relation to the promoter of the present invention include any substitution of, variation of, modification of, replacement of, deletion of or addition of one (or more) nucleic acid from or to the respective promoter sequence providing the resultant promoter sequence allows expression of a GOI, preferably wherein the resultant promoter sequence has at least the same effect as SEQ.I.D. No. 14. In particular, the term "homologue" covers homology with respect to similarity of structure and/or similarity of function providing the resultant promoter sequence has the ability to allow for expression of a GOI, such as a nucleotide sequence according to the present invention. With respect to sequence homology (i.e. similarity), preferably there is more than 80% homology, more preferably at least 85% homology, more preferably at least 90% homology, even more preferably at least 95% homology, more preferably at least 98% homology. The above terms are also synonymous with allelic variations of the sequences.

The intron sequence of the present invention can be any one or all of the intron sequences of the present invention, including partial sequences thereof, provided that if partial sense sequences are used (i.e. sequences that are not or do not comprise any one or more of the full sequences shown as SEQ.I.D. No.1-13) the partial sequences affect enzymatic activity. Suitable examples of partial sequences include sequences that are shorter than any one of the full sense sequences shown as SEQ.I.D.No.s 1 to 13 but which comprise nucleotides that are adjacent the respective exon or exons.

With regard to the second aspect of the present invention (i.e. specifically affecting SBE activity), the nucleotide sequences of the present invention may comprise one or more sense or antisense exon sequences of the SBE gene (but not sense exon sequences naturally associated with the intron sequence), including complete or partial sequences thereof, providing the nucleotide sequences can affect SBE activity, preferably wherein the nucleotide sequences reduce or eliminate SBE activity. Preferably, the nucleotide sequence of the second aspect of the present invention does not comprise sense exon sequences.

The term "vector" includes an expression vector and a transformation vector. The term "expression vector" means a construct capable of *in vivo* or *in vitro* expression. The term "transformation vector" means a construct capable of being transferred from one species to another - such as from an *E. Coli* plasmid to a fungus or a plant cell, or from an *Agrobacterium* to a plant cell.

15

20

25

30

10

5

The term "construct" - which is synonymous with terms such as "conjugate", "cassette" and "hybrid" - in relation to the sense nucleotide sequence aspect of the present invention includes the nucleotide sequence according to the present invention directly or indirectly attached to a promoter. An example of an indirect attachment is the provision of a suitable spacer group such as an intron sequence, such as the Sh1-intron or the ADH intron, intermediate the promoter and the nucleotide sequence of the present invention. The same is true for the term "fused" in relation to the present invention which includes direct or indirect attachment. The terms do not cover the natural combination of the wild type SBE gene when associated with the wild type SBE gene promoter in their natural environment.

The construct may even contain or express a marker which allows for the selection of the genetic construct in, for example, a plant cell into which it has been transferred. Various markers exist which may be used in, for example, plants - such as mannose. Other examples of markers include those that provide for antibiotic resistance - e.g. resistance to G418, hygromycin, bleomycin, kanamycin and gentamycin.

10

15

20

25

30

The construct of the present invention preferably comprises a promoter. The term "promoter" is used in the normal sense of the art, e.g. an RNA polymerase binding site in the Jacob-Monod theory of gene expression. Examples of suitable promoters are those that can direct efficient expression of the nucleotide sequence of the present invention and/or in a specific type of cell. Some examples of tissue specific promoters are disclosed in WO 92/11375.

The promoter could additionally include conserved regions such as a Pribnow Box or a TATA box. The promoters may even contain other sequences to affect (such as to maintain, enhance, decrease) the levels of expression of the nucleotide sequence of the present invention. Suitable examples of such sequences include the *Sh1*-intron or an ADH intron. Other sequences include inducible elements - such as temperature, chemical, light or stress inducible elements. Also, suitable elements to enhance transcription or translation may be present. An example of the latter element is the TMV 5' leader sequence (see Sleat Gene 217 [1987] 217-225; and Dawson Plant Mol. Biol. 23 [1993] 97).

As mentioned, the construct and/or the vector of the present invention may include a transcriptional initiation region which may provide for regulated or constitutive expression. Any suitable promoter may be used for the transcriptional initiation region, such as a tissue specific promoter. In one aspect, preferably the promoter is the patatin promoter or the E35S promoter. In another aspect, preferably the promoter is the SBE promoter.

If, for example, the organism is a plant then the promoter can be one that affects expression of the nucleotide sequence in any one or more of seed, tuber, stem, sprout, root and leaf tissues, preferably tuber. By way of example, the promoter for the nucleotide sequence of the present invention can be the α -Amy 1 promoter (otherwise known as the Amy 1 promoter, the Amy 637 promoter or the α -Amy 637 promoter) as described in our co-pending UK patent application No. 9421292.5 filed 21 October 1994. Alternatively, the promoter for the nucleotide sequence of the present invention can be the α -Amy 3 promoter (otherwise known as the Amy 3

10

15

promoter, the Amy 351 promoter or the α -Amy 351 promoter) as described in our co-pending UK patent application No. 9421286.7 filed 21 October 1994.

The present invention also encompasses the use of a promoter to express a nucleotide sequence according to the present invention, wherein a part of the promoter is inactivated but wherein the promoter can still function as a promoter. Partial inactivation of a promoter in some instances is advantageous. In particular, with the Amy 351 promoter mentioned earlier it is possible to inactivate a part of it so that the partially inactivated promoter expresses the nucleotide sequence of the present invention in a more specific manner such as in just one specific tissue type or organ. The term "inactivated" means partial inactivation in the sense that the expression pattern of the promoter is modified but wherein the partially inactivated promoter still functions as a promoter. However, as mentioned above, the modified promoter is capable of expressing a gene coding for the enzyme of the present invention in at least one (but not all) specific tissue of the original promoter. Examples of partial inactivation include altering the folding pattern of the promoter sequence, or binding species to parts of the nucleotide sequence, so that a part of the nucleotide sequence is not recognised by, for example, RNA polymerase. Another, and preferable, way of partially inactivating the promoter is to truncate it to form fragments thereof. Another way would be to mutate at least a part of the sequence so that the RNA polymerase can not bind to that part or another part. Another modification is to mutate the binding sites for regulatory proteins for example the CreA protein known from filamentous fungi to exert carbon catabolite repression, and thus abolish the catabolite repression of the native promoter.

25

30

20

The construct and/or the vector of the present invention may include a transcriptional termination region.

The nucleotide according to the present invention can be expressed in combination (but not necessarily at the same time) with an additional construct. Thus the present invention also provides a combination of constructs comprising a first construct comprising the nucleotide sequence according to the present invention operatively

linked to a first promoter; and a second construct comprising a GOI operatively linked to a second promoter (which need not be the same as the first promoter). With this aspect of the present invention the combination of constructs may be present in the same vector, plasmid, cells, tissue, organ or organism. This aspect of the present invention also covers methods of expressing the same, preferably in specific cells or tissues, such as expression in just a specific cell or tissue, of an organism, typically a plant. With this aspect of the present invention the second construct does not cover the natural combination of the gene coding for an enzyme ordinarily associated with the wild type gene promoter when they are both in their natural environment.

10

An example of a suitable combination would be a first construct comprising the nucleotide sequence of the present invention and a promoter, such as the promoter of the present invention, and a second construct comprising a promoter, such as the promoter of the present invention, and a GOI wherein the GOI codes for another starch branching enzyme either in sense or antisense orientation.

15

The above comments relating to the term "construct" for the sense nucleotide aspect of the present invention are equally applicable to the term "construct" for the promoter aspect of the present invention. In this regard, the term includes the promoter according to the present invention directly or indirectly attached to a GOI.

20

The term "GOI" with reference to the promoter aspect of the present invention or the combination aspect of the present invention means any gene of interest, which need not necessarily code for a protein or an enzyme - as is explained later. A GOI can be any nucleotide sequence that is either foreign or natural to the organism in question, for example a plant.

25

30

Typical examples of a GOI include genes encoding for other proteins or enzymes that modify metabolic and catabolic processes. The GOI may code for an agent for introducing or increasing pathogen resistance.

10

15

20

25

The GOI may even be an antisense construct for modifying the expression of natural transcripts present in the relevant tissues. An example of such a GOI is the nucleotide sequence according to the present invention.

The GOI may even code for a protein that is non-natural to the host organism - e.g. a plant. The GOI may code for a compound that is of benefit to animals or humans. For example, the GOI could code for a pharmaceutically active protein or enzyme such as any one of the therapeutic compounds insulin, interferon, human serum albumin, human growth factor and blood clotting factors. The GOI may even code for a protein giving additional nutritional value to a food or feed or crop. Typical examples include plant proteins that can inhibit the formation of anti-nutritive factors and plant proteins that have a more desirable amino acid composition (e.g. a higher lysine content than a non-transgenic plant). The GOI may even code for an enzyme that can be used in food processing such as xylanases and α -galactosidase. The GOI can be a gene encoding for any one of a pest toxin, an antisense transcript such as that for α -amylase, a protease or a glucanase. Alternatively, the GOI can be a nucleotide sequence according to the present invention.

The GOI can be the nucleotide sequence coding for the arabinofuranosidase enzyme which is the subject of our co-pending UK patent application 9505479.7. The GOI can be the nucleotide sequence coding for the glucanase enzyme which is the subject of our co-pending UK patent application 9505475.5. The GOI can be the nucleotide sequence coding for the α -amylase enzyme which is the subject of our co-pending UK patent application 9413439.2. The GOI can be the nucleotide sequence coding for the α -amylase enzyme which is the subject of our co-pending UK patent application 9421290.9. The GOI can be any of the nucleotide sequences coding for the α -glucan lyase enzyme which are described in our co-pending PCT patent application PCT/EP94/03397.

In one aspect the GOI can even be a nucleotide sequence according to the present invention but when operatively linked to a different promoter.

The GOI could include a sequence that codes for one or more of a xylanase, an arabinase, an acetyl esterase, a rhamnogalacturonase, a glucanase, a pectinase, a branching enzyme or another carbohydrate modifying enzyme or proteinase. Alternatively, the GOI may be a sequence that is antisense to any of those sequences.

5

As mentioned above, the present invention provides a mechanism for selectively affecting a particular enzymatic activity.

10

In an important application of the present invention it is now possible to reduce or eliminate expression of a genomic nucleotide sequence coding for a genomic protein or enzyme by expressing a sense intron construct for that particular genomic protein or enzyme and (e.g. at the same time) expressing a recombinant version of that enzyme or protein - in other words the GOI is a recombinant nucleotide sequence coding for the genomic enzyme or protein. This application allows expression of desired recombinant enzymes and proteins in the absence of (or reduced levels of) respective genomic enzymes and proteins. Thus the desired recombinant enzymes and proteins can be easily separated and purified from the host organism. This particular aspect of the present invention is very advantageous over the prior art methods which, for example, rely on the use of anti-sense exon expression which methods also affect expression of the recombinant enzyme.

20

25

30

15

Thus, a further aspect of the present invention relates to a method of expressing a recombinant protein or enzyme in a host organism comprising expressing a nucleotide sequence coding for the recombinant protein or enzyme; and expressing a further nucleotide sequence wherein the further nucleotide sequence codes, partially or completely, for an intron in a sense orientation; wherein the intron is an intron normally associated with the genomic gene encoding a protein or an enzyme corresponding to the recombinant protein or enzyme; and wherein the further nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron. Additional aspects cover the combination of those nucleotide sequences including their incorporation in constructs, vectors, cells, tissues and transgenic organisms.

Therefore the present invention also relates to a combination of nucleotide sequences comprising a first nucleotide sequence coding for a recombinant enzyme; and a second nucleotide sequence which corresponds to an intron in a sense orientation; wherein the intron is an intron that is associated with a genomic gene encoding the enzyme corresponding to the recombinant enzyme; and wherein the second nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron.

The GOI may even code for one or more introns but in an antisense orientation, such as any one or more of the antisense intron sequences presented in the attached sequence listings. For example, the present invention also covers the expression of for example a sense intron (e.g. SEQ.I.D.No. 1) in combination with for example an antisense sense intron which preferably is not complementary to the sense intron sequence (e.g. SEQ.I.D.No. 16).

15

10

5

The terms "cell", "tissue" and "organ" include cell, tissue and organ per se and when within an organism.

20

The term "organism" in relation to the present invention includes any organism that could comprise the nucleotide sequence according to the present invention and/or wherein the nucleotide sequence according to the present invention can be expressed when present in the organism. Preferably the organism is a starch producing organism such as any one of a plant, algae, fungi, yeast and bacteria, as well as cell lines thereof. Preferably the organism is a plant.

25

The term "starch producing organism" includes any organism that can biosynthesise starch. Preferably, the starch producing organism is a plant.

30

The term "plant" as used herein includes any suitable angiosperm, gymnosperm, monocotyledon and dicotyledon. Typical examples of suitable plants include vegetables such as potatoes; cereals such as wheat, maize, and barley; fruit; trees; flowers; and other plant crops. Preferably, the term means "potato".

The term "transgenic organism" in relation to the present invention includes any organism that comprises the nucleotide sequence according to the present invention and/or products obtained therefrom, and/or wherein the nucleotide sequence according to the present invention can be expressed within the organism. Preferably the nucleotide sequence of the present invention is incorporated in the genome of the organism. Preferably the transgenic organism is a plant, more preferably a potato.

To prepare the host organism one can use prokaryotic or eukaryotic organisms. Examples of suitable prokaryotic hosts include *E. coli* and *Bacillus subtilis*. Teachings on the transformation of prokaryotic hosts is well documented in the art, for example see Sambrook *et al* (Sambrook *et al*. in Molecular Cloning: A Laboratory Manual, 2nd edition, 1989, Cold Spring Harbor Laboratory Press).

Even though the enzyme according to the present invention and the nucleotide sequence coding for same are not disclosed in EP-B-0470145 and CA-A-2006454, those two documents do provide some useful background commentary on the types of techniques that may be employed to prepare transgenic plants according to the present invention. Some of these background teachings are now included in the following commentary.

20

5

10

15

The basic principle in the construction of genetically modified plants is to insert genetic information in the plant genome so as to obtain a stable maintenance of the inserted genetic material.

25

Several techniques exist for inserting the genetic information, the two main principles being direct introduction of the genetic information and introduction of the genetic information by use of a vector system. A review of the general techniques may be found in articles by Potrykus (Annu Rev Plant Physiol Plant Mol Biol [1991] 42:205-225) and Christou (Agro-Food-Industry Hi-Tech March/April 1994 17-27).

30

Thus, in one aspect, the present invention relates to a vector system which carries a nucleotide sequence or construct according to the present invention and which is

10

15

20

25

capable of introducing the nucleotide sequence or construct into the genome of an organism, such as a plant.

The vector system may comprise one vector, but it can comprise two vectors. In the case of two vectors, the vector system is normally referred to as a binary vector system. Binary vector systems are described in further detail in Gynheung An et al. (1980), Binary Vectors, Plant Molecular Biology Manual A3, 1-19.

One extensively employed system for transformation of plant cells with a given promoter or nucleotide sequence or construct is based on the use of a Ti plasmid from Agrobacterium tumefaciens or a Ri plasmid from Agrobacterium rhizogenes An et al. (1986), Plant Physiol. 81, 301-305 and Butcher D.N. et al. (1980), Tissue Culture Methods for Plant Pathologists, eds.: D.S. Ingrams and J.P. Helgeson, 203-208. Several different Ti and Ri plasmids have been constructed which are suitable for the construction of the plant or plant cell constructs described above. A non-limiting example of such a Ti plasmid is pGV3850.

The nucleotide sequence or construct of the present invention should preferably be inserted into the Ti-plasmid between the terminal sequences of the T-DNA or adjacent a T-DNA sequence so as to avoid disruption of the sequences immediately surrounding the T-DNA borders, as at least one of these regions appears to be essential for insertion of modified T-DNA into the plant genome.

As will be understood from the above explanation, if the organism is a plant the vector system of the present invention is preferably one which contains the sequences necessary to infect the plant (e.g. the *vir* region) and at least one border part of a T-DNA sequence, the border part being located on the same vector as the genetic construct.

Furthermore, the vector system is preferably an Agrobacterium tumefaciens Tiplasmid or an Agrobacterium rhizogenes Ri-plasmid or a derivative thereof. As these plasmids are well-known and widely employed in the construction of transgenic plants, many vector systems exist which are based on these plasmids or derivatives thereof.

In the construction of a transgenic plant the nucleotide sequence or construct of the present invention may be first constructed in a microorganism in which the vector can replicate and which is easy to manipulate before insertion into the plant. An example of a useful microorganism is *E. coli*, but other microorganisms having the above properties may be used. When a vector of a vector system as defined above has been constructed in *E. coli*, it is transferred, if necessary, into a suitable *Agrobacterium* strain, e.g. *Agrobacterium tumefaciens*. The Ti-plasmid harbouring the nucleotide sequence or construct of the present invention is thus preferably transferred into a suitable *Agrobacterium* strain, e.g. *A. tumefaciens*, so as to obtain an *Agrobacterium* cell harbouring the promoter or nucleotide sequence or construct of the present invention, which DNA is subsequently transferred into the plant cell to be modified.

15

20

10

5

If, for example, for the transformation the Ti- or Ri-plasmid of the plant cells is used, at least the right boundary and often however the right and the left boundary of the Ti- and Ri-plasmid T-DNA, as flanking areas of the introduced genes, can be connected. The use of T-DNA for the transformation of plant cells has been intensively studied and is described in EP-A-120516; Hoekema, in: The Binary Plant Vector System Offset-drukkerij Kanters B.B., Alblasserdam, 1985, Chapter V; Fraley, et al., Crit. Rev. Plant Sci., 4:1-46; and An et al., EMBO J. (1985) 4:277-284.

25

30

Direct infection of plant tissues by Agrobacterium is a simple technique which has been widely employed and which is described in Butcher D.N. et al. (1980), Tissue Culture Methods for Plant Pathologists, eds.: D.S. Ingrams and J.P. Helgeson, 203-208. For further teachings on this topic see Potrykus (Annu Rev Plant Physiol Plant Mol Biol [1991] 42:205-225) and Christou (Agro-Food-Industry Hi-Tech March/April 1994 17-27). With this technique, infection of a plant may be performed in or on a certain part or tissue of the plant, i.e. on a part of a leaf, a root, a stem or another part of the plant.

10

15

25

30

Typically, with direct infection of plant tissues by Agrobacterium carrying the GOI (such as the nucleotide sequence according to the present invention) and, optionally, a promoter, a plant to be infected is wounded, e.g. by cutting the plant with a razor blade or puncturing the plant with a needle or rubbing the plant with an abrasive. The wound is then inoculated with the Agrobacterium. The inoculated plant or plant part is then grown on a suitable culture medium and allowed to develop into mature plants.

When plant cells are constructed, these cells may be grown and maintained in accordance with well-known tissue culturing methods such as by culturing the cells in a suitable culture medium supplied with the necessary growth factors such as amino acids, plant hormones, vitamins, etc.

Regeneration of the transformed cells into genetically modified plants may be accomplished using known methods for the regeneration of plants from cell or tissue cultures, for example by selecting transformed shoots using an antibiotic and by subculturing the shoots on a medium containing the appropriate nutrients, plant hormones, etc.

Further teachings on plant transformation may be found in EP-A-0449375.

As reported in CA-A-2006454, a large amount of cloning vectors are available which contain a replication system in *E. coli* and a marker which allows a selection of the transformed cells. The vectors contain for example pBR 322, pUC series, M13 mp series, pACYC 184 etc. In this way, the nucleotide or construct of the present invention can be introduced into a suitable restriction position in the vector. The contained plasmid is then used for the transformation in *E. coli*. The *E. coli* cells are cultivated in a suitable nutrient medium and then harvested and lysed. The plasmid is then recovered. As a method of analysis there is generally used sequence analysis, restriction analysis, electrophoresis and further biochemical-molecular biological methods. After each manipulation, the used DNA sequence can be restricted and connected with the next DNA sequence. Each sequence can be cloned in the same

20

or different plasmid.

After the introduction of the nucleotide sequence or construct according to the present invention in the plants the presence and/or insertion of further DNA sequences may be necessary - such as to create combination systems as outlined above (e.g. an organism comprising a combination of constructs).

The above commentary for the transformation of prokaryotic organisms and plants with the nucleotide sequence of the present invention is equally applicable for the transformation of those organisms with the promoter of the present invention.

In summation, the present invention relates to affecting enzyme activity by expressing sense intron sequences.

Also, the present invention relates to a promoter useful for the expression of those sense intron sequences.

The following samples have been deposited in accordance with the Budapest Treaty at the recognised depositary The National Collections of Industrial and Marine Bacteria Limited (NCIMB) at 23 St Machar Drive, Aberdeen, Scotland. AB2 1RY, United Kingdom, on 13 July 1995:

NCIMB 40754 (which refers to pBEA 11 as described herein);

NCIMB 40751 (which refers to λ -SBE 3.2 as described herein), and

NCIMB 40752 (which refers to λ -SBE 3.4 as described herein).

A highly preferred embodiment of the present invention therefore relates to a method of affecting enzymatic activity in a plant (or a cell, a tissue or an organ thereof) comprising expressing in the plant (or a cell, a tissue or an organ thereof) a nucleotide sequence wherein the nucleotide sequence codes, partially or completely,

for an intron in a sense orientation; wherein the nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron; wherein starch branching enzyme activity is affected and/or the levels of amylopectin are affected and/or the composition of starch is changed; and wherein the intron nucleotide sequence is obtainable from NCIMB 40751, NCIMB 40752, or NCIMB 40754 or a variant, derivative or homologue thereof.

A more highly preferred aspect of the present invention therefore relates to a method of affecting enzymatic activity in a plant (or a cell, a tissue or an organ thereof) comprising expressing in the plant (or a cell, a tissue or an organ thereof) a nucleotide sequence wherein the nucleotide sequence codes, partially or completely, for an intron in a sense orientation; wherein the nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron; wherein starch branching enzyme activity is affected and/or the levels of amylopectin are affected and/or the composition of starch is changed; wherein the nucleotide sequence comprises the sequence shown as any one of SEQ.I.D. No. 1 to SEQ.I.D. No. 13 or a variant, derivative or homologue thereof, including combinations thereof; and wherein the intron nucleotide sequence is obtainable from NCIMB 40751, NCIMB 40752, or NCIMB 40754, or a variant, derivative or homologue thereof.

20

25

5

10

15

The present invention will now be described only by way of example, in which reference is made to the following attached Figures:

Figure 1, which is a schematic representation of the biosynthesis of amylose and amylopectin;

- Figure 2, which is a diagrammatic representation of the α -1-4-links and the α -1-6 links of amylopectin;
- Figure 3, which is a diagrammatic representation of the exon-intron structure of a genomic SBE clone;

- Figure 4, which is a plasmid map of pPATA1, which is 3936 bp in size;
- Figure 5, which is a plasmid map of pABE7, which is 5106 bp in size;
- Figure 6, which is a plasmid map of pVictorIV Man, which is 7080 bp in size;
 - Figure 7, which is a plasmid map of pBEA11, which is 9.54 kb in size;
- Figure 8, which shows the full genomic nucleotide sequence for SBE including the promoter, exons and introns;
 - Figure 9, which is a plasmid map of pVictor5a, which is 9.12 kb in size; and
 - Figure 10, which is a plasmid map of pBEP2, which is 10.32 kb in size.

20

letters.

Figures 1 and 2 were referred to above in the introductory description concerning starch in general. As mentioned, Figure 3 is a diagrammatic representation of the exon-intron structure of a genomic SBE clone, the sequence of which is shown in Figure 8. This clone, which has about 11.5 k base pairs, comprises 14 exons and 13 introns. The introns are numbered in increasing order from the 5' end to the 3' end and correspond to SEQ.I.D.No.s 1-13, respectively. Their respective antisense intron sequences are shown as SEQ.I.D.No.s 15-27.

In more detail, Figures 3 and 8 present information on the 11468 base pairs of a potato SBE gene. The 5' region from nucleotides 1 to 2082 contain the promoter region of the SBE gene. A TATA box candidate at nucleotide 2048 to 2051 is boxed. The homology between a potato SBE cDNA clone (Poulsen & Kreiberg (1993) Plant Physiol 102: 1053-1054) and the exon DNAs begin at 2083 bp and end at 9666 bp. The homology between the cDNA and the exon DNA is indicated by nucleotides in upper case letters, while the translated amino acid sequences are shown in the single letter code below the exon DNA. Intron sequences are indicated by lower case

Figure 7 is a plasmid map of pBEA7, which is 9.54 k base pairs in size. Plasmid pBEA 11 comprises the first intron sequence of the potato SBE gene. This first intron sequence, which has 1177 base pairs, is shown in Figure 3 and lies between the first exon and the second exon.

5

These experiments and aspects of the present invention are now discussed in more detail.

EXPERIMENTAL PROTOCOL

10

ISOLATION, SUBCLONING IN PLASMIDS, AND SEQUENCING OF GENOMIC SBE CLONES

15

20

Various clones containing the potato SBE gene were isolated from a Desiree potato genomic library (Clontech Laboratories Inc., Palo Alto CA, USA) using radioactively labelled potato SBE cDNA (Poulsen & Kreiberg (1993) Plant Physiol. 102:1053-1054) as probe. The fragments of the isolated λ-phages containing SBE DNA (λSBE 3.2 - NCIMB 40751 - and λSBE-3.4 - NCIMB 40752) were identified by Southern analysis and then subcloned into pBluescript II vectors (Clontech Laboratories Inc., Palo Alto CA, USA). λSBE 3.2 contains a 15 kb potato DNA insert and λSBE-3.4 contains a 13 kb potato DNA insert. The resultant plasmids were called pGB3, pGB11, pGB15, pGB16 and pGB25 (see discussion below). The respective inserts were then sequenced using the Pharmacia Autoread Sequencing Kit (Pharmacia, Uppsala) and a A.L.F. DNA sequencer (Pharmacia, Uppsala).

25

30

In total, a stretch of 11.5 kb of the SBE gene was sequenced. The sequence was deduced from the above-mentioned plasmids, wherein: pGB25 contains the sequences from 1 bp to 836 bp, pGB15 contains the sequences from 735 bp to 2580 bp, pGB16 contains the sequences from 2580 bp to 5093 bp, pGB11 contains the sequences from 3348 bp to 7975 bp, and pGB3 contains the sequences from 7533 bp to 11468 bp.

In more detail, pGB3 was constructed by insertion of a 4 kb EcoRI fragment isolated from λSBE 3.2 into the EcoRI site of pBluescript II SK (+). pGB11 was constructed by insertion of a 4.7 kb XhoI fragment isolated from λSBE 3.4 into the XhoI site of pBluescript II SK (+). pGB15 was constructed by insertion of a 1.7 kb SpeI fragment isolated from λSBE 3.4 into the SpeI site of pBluescript II SK (+). pGB16 was constructed by insertion of a 2.5 kb SpeI fragment isolated from λSBE 3.4 into the SpeI site of pBluescript II SK (+). For the construction of pGB25 a PCR fragment was produced with the primers

10 5' GGA ATT CCA GTC GCA GTC TAC ATT AC 3'

and

5' CGG GAT CCA GAG GCA TTA AGA TTT CTG G 3'

15 ·

5

and λ SBE 3.4 as a template.

The PCR fragment was digested with *BamH*I and *EcoR*I, and inserted in pBluescript II SK (+) digested with the same restriction enzymes.

20

CONSTRUCTION OF PLASMID pBEA11

The SBE intron 1 was amplified by PCR using the oligonucleotides

5' CGG GAT CCA AAG AAA TTC TCG AGG TTA CAT GG 3'

and

5' CGG GAT CCG GGG TAA TTT TTA CTA ATT TCA TG 3'

30

25

and the λ SBE 3.4 phage containing the SBE gene as template.

The PCR product was digested with *BamH*I and inserted in a sense orientation in the *BamH*I site of plasmid pPATA1 (described in WO 94/24292) between the patatin promoter and the 35S terminator. This construction, pABE7, was digested with *Kpn*I, and the 2.4 kb "patatin promoter-SBE intron 1-35S terminator" *Kpn*I fragment was isolated and inserted in the *Kpn*I site of the plant transformation vector pVictorIV Man yielding plasmid pBEA11.

PRODUCTION OF TRANSGENIC POTATO PLANTS

10 Axenic stock cultures

Shoot cultures of *Solanum tuberosum* 'Bintje' and 'Dianella' are maintained on a substrate (LS) of a formula according to Linsmaier, E.U. and Skoog, F. (1965), Physiol. Plant. 18: 100-127, in addition containing 2 μ M silver thiosulphate at 25°C and 16 h light/8 h dark.

The cultures were subcultured after approximately 40 days. Leaves were then cut off the shoots and cut into nodal segments (approximately 0.8 cm) each containing one node.

20

25

30

15

5

Inoculation of potato tissues

Shoots from approximately 40 days old shoot cultures (height approximately 5-6 cms) were cut into internodal segments (approximately 0.8 cm). The segments were placed into liquid LS-substrate containing the transformed Agrobacterium tumefaciens containing the binary vector of interest. The Agrobacterium were grown overnight in YMB-substrate (di-potassium hydrogen phosphate, trihydrate (0.66 g/l); magnesium sulphate, heptahydrate (0.20 g/l); sodium chloride (0.10 g/l); mannitol (10.0 g/l); and yeast extract (0.40 g/l)) containing appropriate antibiotics (corresponding to the resistance gene of the Agrobacterium strain) to an optical density at 660 nm (OD-660) of approximately 0.8, centrifuged and resuspended in the LS-substrate to an OD-660 of 0.5.

The segments were left in the suspension of Agrobacterium for 30 minutes and then the excess of bacteria were removed by blotting the segments on sterile filter paper.

Co-cultivation

5

The shoot segments were co-cultured with bacteria for 48 hours directly on LS-substrate containing agar (8.0 g/l), 2,4-dichlorophenoxyacetic acid (2.0 mg/l) and trans-zeatin (0.5 mg/l). The substrate and also the explants were covered with sterile filter papers, and the petri dishes were placed at 25°C and 16 h light/8 dark.

10

15

"Washing" procedure

After the 48 h on the co-cultivation substrate the segments were transferred to containers containing liquid LS-substrate containing 800 mg/l carbenicillin. The containers were gently shaken and by this procedure the major part of the *Agrobacterium* was either washed off the segments and/or killed.

Selection

20

25

30

After the washing procedure the segments were transferred to plates containing the LS-substrate, agar (8 g/l), trans-zeatin (1-5 mg/l), gibberellic acid (0.1 mg/l), carbenicillin (800 mg/l), and kanamycin sulphate (50-100 mg/l) or phosphinotricin (1-5 mg/l) or mannose (5 g/l) depending on the vector construction used. The segments were sub-cultured to fresh substrate each 3-4 weeks. In 3 to 4 weeks, shoots develop from the segments and the formation of new shoots continued for 3-4 months.

Rooting of regenerated shoots

The regenerated shoots were transferred to rooting substrate composed of LS-substrate, agar (8 g/l) and carbenicillin (800 mg/l).

The transgenic genotype of the regenerated shoot were verified by testing the rooting ability on the above mentioned substrates containing kanamycin sulphate (200 mg/l), by performing NPTII assays (Radke, S. E. et al, Theor. Appl. Genet. (1988), 75: 685-694) or by performing PCR analysis according to Wang et al (1993, NAR $\underline{21}$ pp 4153-4154). Plants which were not positive in any of these assays were discarded or used as controls. Alternatively, the transgenic plants could be verified by performing a GUS assay on the co-introduced β -glucuronidase gene according to Hodal, L. et al. (Pl. Sci. (1992), 87: 115-122).

10 Transfer to soil

5

15

25

The newly rooted plants (height approx. 2-3 cms) were transplanted from rooting substrate to soil and placed in a growth chamber (21°C, 16 hour light 200-400uE/m²/sec). When the plants were well established they were transferred to the greenhouse, where they were grown until tubers had developed and the upper part of the plants were senescing.

Harvesting

The potatoes were harvested after about 3 months and then analysed.

BRANCHING ENZYME ANALYSIS

The SBE expression in the transgenic potato lines were measured using the SBE assays described by Blennow and Johansson (Phytochemistry (1991) 30:437-444) and by standard Western procedures using antibodies directed against potato SBE.

STARCH ANALYSIS

30 Starch was isolated from potato tubers and analysed for the amylose:amylopectin ratio (Hovenkamp-Hermelink et al. (1988) Potato Research 31:241-246). In addition, the chain length distribution of amylopectin was determined by analysis of isoamylase

digested starch on a Dionex HPAEC. The number of reducing ends in isoamylase digested starch was determined by the method described by N. Nelson (1944) J. Biol.Chem. 153:375-380.

The results revealed that there was a reduction in the level of synthesis of SBE and/or the level of activity of SBE and/or the composition of starch SBE in the transgenic plants.

CONSTRUCTION OF SBE PROMOTER CONSTRUCT

10

20

An SBE promoter fragment was amplified from λ -SBE 3.4 using primers:

5' CCA TCG ATA CTT TAA GTG ATT TGA TGG C 3'

15 and

5' CGG GAT CCT GTT CTG ATT CTT GAT TTC C 3'.

The PCR product was digested with *ClaI* and *BamHI*. The resultant 1.2 kb fragment was then inserted in pVictor5a (see Figure 9) linearised with *ClaI* and *BgIII* yielding pBEP2 (see Figure 10).

STARCH BRANCHING ENZYME MEASUREMENTS OF POTATO TUBERS

Potatoes from potato plants transformed with pBEA11 were cut in small pieces and homogenised in extraction buffer (50 mM Tris-HCl pH 7.5, Sodium-dithionit (0.1 g/l), and 2 mM DTT) using a Ultra-Turax homogenizer; 1 g of Dowex xl. was added pr. 10 g of tuber. The crude homogenate was filtered through a miracloth filter and centrifuged at 4°C for 10 minutes at 24.700 g. The supernatant was used for starch branching enzyme assays.

PCT/EP96/03053

The starch branching enzyme assays were carried out at 25 oC in a volume of 400 μ l composed of 0.1 M Na citrate buffer pH 7.0, 0.75 mg/ml amylose, 5 mg/ml bovine serum albumin and the potato extract. At 0, 15 30 and 60 minutes aliquots of 50 μ l were removed from the reaction into 20 μ l 3 N HCl. 1 ml of iodine solution was added and the decrease in absorbance at 620 nm was measured with an ELISA spectrophotometer.

The starch branching enzyme (SBE) levels in tuber extracts were measured from 24 transgenic Dianella potato plants transformed with plasmid pBEA11.

10

The results showed that the BEA11 transgenic lines produced tubers which have SBE levels that are only 10 % to 15 % of the SBE levels found in non transformed Dianella plants.

15 SUMMATION

The above-mentioned examples relate to the isolation and sequencing of a gene for potato SBE. The examples further demonstrate that it is possible to prepare SBE intron constructs. These SBE intron constructs can be introduced into plants, such as potato plants. After introduction, a reduction in the level of synthesis of SBE and/or the level of activity of SBE and/or the composition of starch in plants can be achieved.

25

30

20

Without wishing to be bound by theory it is believed that the expressed sense intron nucleotide sequence according to the present invention affects enzymatic activity via co-suppression and/or trans-activation. Reviews of these mechanisms has been published by Finnegan and McElroy (1994 Biotechnology 12 pp 883 - 887) and Matzke and Matzke (1995 TIG 11 No. 1 pp 1 - 3). By these mechanisms, it is believed that the sense introns of the present invention reduce the level of plant enzyme activity (in particular SBE activity), which in turn for SBE activity is believed to influence the amylose:amylopectin ratio and thus the branching pattern of amylopectin.

15

20

Thus, the present invention provides a method wherein it is possible to manipulate the starch composition in plants, or tissues or cells thereof, such as potato tubers, by reducing the level of SBE activity by using sense intron sequences.

In summation the present invention therefore relates to the surprising use of sense intron sequences in a method to affect enzymatic activity in plants.

Other modifications of the present invention will be apparent to those skilled in the art without departing from the scope of the present invention. For example, it may be possible to use antisense promoter sequences to affect enzymatic activity, such as antisense SBE promoter - such as a nucleotide sequence comprising the nucleotide sequence shown as SEQ. I.D. No. 28 or a variant, derivative or homologue thereof.

The following pages present a number of sequence listings which have been consecutively numbered from SEQ.I.D. No. 1 - SEQ.I.D. No. 29. In brief, SEQ.I.D. No. 1 - SEQ.I.D. No. 13 represent sense intron sequences (genomic DNA); SEQ.I.D. No. 14 represents the SBE promoter sequence (genomic sequence); SEQ.I.D. No. 15 - SEQ.I.D. No. 27 represent antisense intron sequences; and SEQ. I.D. No. 28 represents the sequence complementary to the SBE promoter sequence i.e. the SBE promoter sequence in antisense orientation. The full genomic nucleotide sequence for SBE including the promoter, exons and introns is shown as SEQ. I.D. No. 29 (see Figures 3 and 8 which highlight particular gene features).

SEQUENCE INFORMATION

SEQ.I.D. No. 1
Intron 1 sequence (1167 bp).

GTAATTTTACTAATTTCATGTTAATTTCAATTATTTTTAGCCTTTGCATTTCCAATATATCT **AGCCATATGTAAGTATCATGTGACAAATTTGCAAGGTGGTTGAGTGTATAAAATTCAAAAATTGAGAGA** TGGAGGGGGGGGGBARAGACAATATTTAGAAAGAGTGTTCTAGGAGGTTATGGAGGACACGGATG AGGGGTAGAAGGTTAGTTAGGTATTTGAGTGTTGTCTGGCTTATCCTTTCATACTAGTAGTCGTGGAAT ATTGTATTATATATCTTGTCGTAGTTATTGTTCCTCGGTAAGAATGCTCTAGCATGCTTCCTTTAGTGT TTTATCATGCCTTCTTTATATTCGCGTTGCTTTGAAATGCTTTTACTTTAGCCGAGGGTCTATTAGAAA CAATCTCTCTATCTCGTAAGGTAGGGGTAAAGTCCTCACCACACTCCACTTGTGGGATTACATTGTGTT TGTTGTTGTAAATCAATTATGTATACATAATAAGTGGATTTTTTACAACACAAATACATGGTCAAGGGC AAAGTTCTGAACACATAAAGGGTTCATTATATGTCCAGGGATATGATAAAAATTGTTTCTTTGTGAAAG TTATATAAGATTTGTTATGGCTTTTGCTGGAAACATAATAAGTTATAATGCTGAGATAGCTACTGAAGT TTGTTTTTCTAGCCTTTTAAATGTACCAATAATAGATTCCGTATCGAACGAGTATGTTTTGATTACCT GGTCATGATGTTTCTATTTTTTACATTTTTTTGGTGTTGAACTGCAATTGAAAATGTTGTATCCTATGA GACGGATAGTTGAGAATGTGTTCTTTGTATGGACCTTGAGAAGCTCAAACGCTACTCCAATAATTTCTA TGAATTCAAATTCAGTTTATGGCTACCAGTCAGTCAGAAATTAGGATATGCTGCATATACTTGTTCAA TTATACTGTAAAATTTCTTAAGTTCTCAAGATATCCATGTAACCTCGAGAATTTCTTTGACAG

SEQ.I.D. No. 2 Intron 2 sequence (321 bp).

SEQ.I.D. No. 3
Intron 3 sequence (504 bp).

SEQ.I.D. No. 4

Intron 4 sequence (146 bp).

GTAGGTCCTCGTCTACTACAAAATAGTAGTTTCCATCATCATAACAGATTTTCCTATTAAAGCATGATG
TTGCAGCATCATTGGCTTTCTTACATGTTCTAATTGCTATTAAGGTTATGCTTCTAATTAACTCATCCA
CAATGCAG

SEQ.I.D. No. 5

Intron 5 sequence (218 bp).

GTTTTGTTATTCATACCTTGAAGCTGAATTTTGAACACCATCATCACGGCATTTCGATTCATGTTCTT
ACTAGTCTTGTTATGTAAGACATTTTGAAATGCAAAAGTTAAAATAATTGTGTCTTTACTAATTTGGAC
TTGATCCCATACTCTTTCCCTTAACAAAATGAGTCAATTCTATAAGTGCTTGAGAACTTACTACTACTCAG
CAATTAAACAG

SEQ.I.D. No. 6

Intron 6 sequence (198 bp).

SEQ.I.D. No. 7

Intron 7 sequence (208bp)

GTTTGTCTGTTTCTATTGCATTTTAAGGTTCATATAGGTTAGCCACGGAAAATCTCACTCTTTGTGAGG TAACCAGGGTTCTGATGGATTATTCAATTTTCTCGTTTATCATTTGTTTATTCTTTTCATGCATTGTGT TTCTTTTTCAATATCCCTCTTATTTGGAGGTAATTTTTCTCATCTATTCACTTTTAGCTTCTAACCACAG

SEQ.I.D. No. 8

Intron 8 sequence (293 bp).

SEQ.I.D. No. 9

Intron 9 sequence (376 bp).

GTTCAAGTATTTTGAATCGCAGCTTGTTAAATAATCTAGTAATTTTTAGATTGCTTACTTGGAAGTCTA
CTTGGTTCTGGGGATGATAGCTCATTTCATCTTGTTCTACTTATTTTCCAACCGAATTTCTGATTTTTG
TTTCGAGATCCAAGTATTAGATTCATTTACACTTATTACCGCCTCATTTCTACCACTAAGGCCTTGATG
AGCAGCTTAAGTTGATTCTTTGAAGCTATAGTTTCAGGCTACCAATCCACAGCCTGCTATATTTGTTGG

ATACTTACCTTTTCTTTACAATGAAGTGATACTAATTGAAATGGTCTAAATCTGATATCTATATTTCTCCCCCTCATGATGAAATGCAG

SEQ.I.D. No. 10

Intron 10 sequence (172 bp).

GTAAAATCATCTAAAGTTGAAAGTGTTGGGTTTATGAAGTGCTTTAATTCTATCCAAGGACAAGTAGAA ACCTTTTTACCTTCCATTTCTTGATGATGGATTTCATATTATTTAATCCAATAGCTGGTCAAATTCGGT AATAGCTGTACTGATTAGTTACTTCACTTTGCAG

SEQ.I.D. No. 11

Intron 11 sequence (145 bp).

GTATATATGTTTTACTTATCCATGAAATTATTGCTCTGCTTGTTTTTAATGTACTGAACAAGTTTTATG
GAGAAGTAACTGAAACAAATCATTTTCACATTGTCTAATTTAACTCTTTTTTCTGATCCTCGCATGACG
AAAACAG

SEQ.I.D. No. 12

Intron 12 sequence (242 bp).

GTAAGGATTTGCTTGAATAACTTTTGATAATAAGATAACAGATGTAGGGTACAGTTCTCTCACCAAAAA GAACTGTAATTGTCTCATCCATCTTTAGTTGTATAAGATATCCGACTGTCTGAGTTCGGAAGTGTTTGA GCCTCCTGCCCTCCCCTGCGTTGTTTAGCTAATTCAAAAAGGAGAAAACTGTTTATTGATGATCTTTG TCTTCATGACAACAGCTGACAAACAGCTGTTTATTGATGATCTTTG

SEQ.I.D. No. 13

Intron 13 sequence (797 bp).

GTACAGTTCTTGCCGTGTGACCTCCCTTTTTATTGTGGTTTTGTTCATAGTTATTTGAATGCGATAGAA
GTTAACTATTGATTACCGCCACAATCGCCAGTTAAGTCCTCTGAACTACTAATTTGAAAGGTAGGAATA
GCCGTAATAAGGTCTACTTTTGGCATCTTACTGTTACAAAACAAAAGGATGCCAAAAAAATTCTTCTCT
ATCCTCTTTTTCCCTAAACCAGTGCATGTAGCTTGCACCTGCATAAACTTAGGTAAATGATCAAAAATG
AAGTTGATGGGAACTTAAAACCGCCCTGAAGTAAAGCTAGGAATAGTCATATAATGTCCACCTTTGGTG
TCTGCGCTAACATCAACAACAACATACCTCGTGTAGTCCCACAAAGTGGTTTCAGGGGGGAGGGTAGAGT
GTATGCAAAACTTACTCCTATCTCAGAGGTAGAGAGAGTTTTTTCAATAGACCCTTGGCTCAAGAAAA
AAGTCCAAAAAGAAGAACAACAACAACATGACAACATGTGTAGCTAAAGCGACCCAACTTGTTTGGGACT
GAAGTAGTTGTTGTTGTAAACAGTGCATGTAGATGAACACATGTCAGAAAATGGACAACACAGTTAT
TTTGTGCAAGTCAAAAAAAATGTACTACTATTTCTTTGTGCAGCTTTATGTATAGAAAAAGTTAAATAACT
AATGAATTTTGCTAGCAGAAAAAATAGCTTGGAGAGAAATTTTTTTATATTGAACTAAGCTAACTATATTC
ATCTTTCTTTTTTGCTTCTTCTCTCCTTGTTTGTGAAG

SEQ.I.D. No. 14 DNA sequence of the SBE gene promoter region.

ATCATGGCCAATTACTGGTTCAAATGCATTACTTCCTTTCAGATTCTTTCGAGTTCTCAT	60
GACCGGTCCTACTACAGACGATACTAACCCGTGGAACTGTTGCATCTGCTTCTTAGAACT	120
CTATGGCTATTTTCGTTAGCTTGGCGTCGGTTTGAACATAGTTTTTGTTTTCAAACTCTT	180
CATTTACAGTCAAAATGTTGTATGGTTTTTGTTTTCCTCAATGATGTTTACAGTGTTGTG	240
TTGTCATCTGTACTTTTGCCTATTACTTGTTTTGAGTTACATGTTAAAAAAGTGTTTATT	300
TTGCCATATTTTGTTCTCTTATTATTATTATCATACATAC	-360
AGTACACAGATCTTAACGTTTATGTTCAATCAACTTTTGGAGGCATTGACAGGTACCACA	420
AATTTTGAGTTTATGATTAAGTTCAATCTTAGAATATGAATTTAACATCTATTATAGATG	480
CATAAAAATAGCTAATGATAGAACATTGACATTTGGCAGAGCTTAGGGTATGGTATATCC	540
AACGTTAATTTAGTAATTTTTGTTACGTACGTATATGAAATATTGAATTAATCACATGAA	600
CGGTGGATATTATTATGAGTTGGCATCAGCAAAATCATTGGTGTAGTTGACTGTAGTT	660
GCAGATTTAATAATAAAATGGTAATTAACGGTCGATATTAAAATAACTCTCATTTCAAGT	720
GGGATTAGAACTAGTTATTAAAAAAATGTATACTTTAAGTGATTTGATGGCATATAATTT	780
AAAGTTTTTCATTCATGCTAAAATTGTTAATTATTGTAATGTAGACTGCGACTGGAATT	840
ATTATAGTGTAAATTTATGCATTCAGTGTAAAATTAAAGTATTGAACTTGTCTGTTTTAG	900
AAAATACTTTATACTTTAATATAGGATTTTGTCATGCGAATTTAAATTAATCGATATTGA	960
ACACGGAATACCAAAATTAAAAAGGATACACATGGCCTTCATATGAACCGTGAACCTTTG	1020
ATAACGTGGAAGTTCAAAGAAGGTAAAGTTTAAGAATAAACTGACAAATTAATT	1080
ATTTGGCCCACTACTAAATTTGCTTTACTTTCTAACATGTCAAGTTGTGCCCTCTTAGTT	1140
GAATGATATTCATTTTCATCCCATAAGTTCAATTTGATTGTCATACCACCCATGATGTT	1200
CTGAAAAATGCTTGGCCATTCACAAAGTTTATCTTAGTTCCTATGAACTTTATAAGAAGC	1260
TTTAATTTGACATGTTATTTATATTAGATGATATAATCCATGACCCAATAGACAAGTGTA	1320
TTAATATTGTAACTTTGTAATTGAGTGTGTCTACATCTTATTCAATCATTTAAGGTCATT	1380
AAAATAAATTATTTTTTGACATTCTAAAACTTTAAGCAGAATAAATA	1440
TAAAAACAAAAAACGACTTATTTATAAATCAACAAACAATTTTAGATTGCTCCAACATAT	1500
TTTTCCAAATTAAATGCAGAAAATGCATAATTTTATACTTGATCTTTATAGCTTATTTTT	1560
TTTAGCCTAACCAACGAATATTTGTAAACTCACAACTTGATTAAAAGGGATTTACAACAA	1620
GATATATAAAGTAGTGACAAATCTTGATTTTAAATATTTTAATTTGGAGGTCAAAATTT	1680
TACCATAATCATTTGTATTTATAATTAAATTTTAAATATCTTATTTAT	1740
AACTTTTAAATATACGTATATACAAAATATAAAATTATTGGCGTTCATATTAGGTCAATA	1800
AATCCTTAACTATATCTGCCTTACCACTAGGAGAAAGTAAAAAACTCTTTACCAAAAATA	1860
CATGTATTATGTATACAAAAGTCGATTAGATTACCTAAATAGAAATTGTATAACGAGTA	1920
AGTAAGTAGAAATATAAAAAAACTACAATACTAAAAAAAA	1980
AAACTAATGGGGTCTGAGTGAAATATTCAGAAAGGGGAGGACTAACAAAAGGGTCATAAT	2040
GTTTTTTTATAAAAGCCACTAAAATGAGGAAATCAAGAATCAGAACATACAAGAAGGCA	2100
GCAGCTGAAGCAAAGTACCATAATTTAATCAATGGAAATTAATT	2160
ACCCATTCG	

SEQ.I.D. No. 15

Intron 1 antisense sequence (1167 bp).

CTGTCAAAGAAATTCTCGAGGTTACATGGATATCTTGAGAACTTAAGAAATTTTACAGTATAATTGAAC AAGTATATGCAGCATATCCTAATTTCTGGACTGACTGGTAGCCATAAACTGAATTTGAATTCATAGAAA TTATTGGAGTAGCGTTTGAGCTTCTCAAGGTCCATACAAAGAACACATTCTCAACTATCCGTCTCATAG GATACAACATTTCAATTGCAGTTCAACACCAAAAAAATGTAAAAAATAGAAACATCATGACCAGGTAA GTAGCTATCTCAGCATTATAACTTATTATGTTTCCAGCAAAAGCCATAACAAATCTTATATAACTTTCA CAAAGAAACAATTTTTATCATATCCCTGGACATATAATGAACCCTTTATGTGTTCAGAACTTTGCCCTT ATGTAATCCCACAAGTGGAGTGTGGTGAGGACTTTACCCCTACCTTACGAGATAGAGAGATTGTTTCTA ATAGACCCTCGGCTAAAGTAAAAGCATTTCAAAGCAACGCGAATATAAAGAAGGCATGATAAAACACTA AAGGAAGCATGCTAGAGCATTCTTACCGAGGAACAATAACTACGACAAGATATATAATACAATAATCGA AGTACAAGAAACAGAAAATAGAATAACAAAGATCAAATAACAAAACAAGAAACTACCCAAATAATTCCA TGTCCTCCATAACCTCCTAGAACACTCTTTCTAAATATTGTCTYTVCCCCCACCCCCCCCCCATCTCTC AATTTTTGAATTTTATACACTCAACCACCTTGCAAATTTGTCACATGATACTTACATATGGCTCTACAA GTGTCATTTTCTTCCATATTTGATATTATAAAAAATAAAAATAAAAACTAAGGAGATGATCCAGATAT ATTGGAAAATGAAATGCAAAGGCTAAAAATAATTGAAATTAACATGAAATTAGTAAAAATTAC

SEQ.I.D. No. 16

Intron 2 antisense sequence (321 bp).

MMMVGCAAGCAATGCACCACAGTTAGTTTATATCAAAAAGAAGAAGGTATTAACGGAGCTAAAAACTG
TTATATACCACATGAAAGAAGTTGATAATGTGAAAACACCATGCTCATAAAGATTGTAATTCAAATAAC
AAATGCCCACAGGAGTAAAGAGCTGTCTTTCCCAAGTTAAGGTATTATAAATTGGCGGAACGAAGTAAC
ACATGTTTGACATCTCCACACGGTGCACAGATCAAATATGCCATGAGCACCAGTCCAGAAGTTTTCCAA
CTATTTATATACTATCCATGCAACCATATAAATTATCAAACATAC

SEQ.I.D. No. 17

Intron 3 antisense sequence (504 bp).

SEQ.I.D. No. 18

Intron 4 antisense sequence (146 bp).

CTGCATTGTGGATGAGTTAATTAGAAGCATAACCTTAATAGCAATTAGAACATGTAAGAAAGCCAATGA TGCTGCAACATCATGCTTTAATAGGAAAATCTGTTATGATGATGGAAACTACTATTTTGTAGTAGACGA GGACCTAC

SEQ.I.D. No. 19

Intron 5 antisense sequence (218 bp).

CTGTTTAATTGCTGAAGTAGTAAGTTCTCAAGCACTTATAGAATTGACTCATTTTGTTAAGGGAAAGAG
TATGGGATCAAGTCCAAATTAGTAAAGACACAATTATTTTAACTTTTGCATTTCAAAATGTCTTACATA
ACAAGACTAGTAAGAACATGAATCGAAATGCCTGTGATGATGGTGTTCAAAATTCAGCTTCAAGGTATG
AATAACAAAAC

SEQ.I.D. No. 20

Intron 6 antisense sequence (198 bp).

SEQ.I.D. No. 21

Intron 7 antisense sequence (208 bp).

CTGTGGTTAGAAGCTAAAAGTGAATAGATGAGAAAAATTACCTCCAAATAAGAGGGATATTGAAAAAGA AACACAATGCATGAAAAGAATAAACAAATGATAAACGAGAAAATTGAATAATCCATCAGAACCCTGGTT ACCTCACAAAGAGTGAGATTTTCCGTGGCTAACCTATATGAACCTTAAAATGCAATAGAAACAGACAAAC

SEQ.I.D. No. 22

Intron 8 antisense sequence (293 bp).

CTGTACAAGTTCATCAAACATTTCACAATTACTCCAAAACAGACACTTGCAAACTCTATACAGTAAT
CTTCTATACTACAAAAAAGTAAACAATGTTTTTTTTAAGATGACATTTGTTCTCAGCAACATAATAGAA
ATCCCTAGACAATGGAAACATTCATCATGTTGTTTTCCTCTATGTTTCAACCCCTTTGATGTTCAACAG
TTCAGGTCATTTTGAGGAATGAATCTTGTTCAAGTAAGCCAAACTAATTGTAATTATCACAAAATATCT
AAAGATGTAAGACATAC

SEQ.I.D. No. 23

Intron 9 antisense sequence (376 bp).

CTGCATTCATGAGGGGGGGGAGAAAGACGGAGAAATATAGATATCAGATTTAGACCATTTCAATTAG
TATCACTTCATTGTAAAGAAAAGGTAAGTATCCAACAAATATAGCAGGCTGTGGATTGGTAGCCTGAAA
CTATAGCTTCAAAGAATCAACTTAAGCTGCTCATCAAGGCCTTAGTGGTAGAAATGAGGCGGTAATAAG
TGTAAATGAATCTAATACTTGGATCTCGAAACAAAAATCAGAAATTCGGTTGGAAAATAAGTAGAACAA

GATGAAATGAGCTATCATCCCCAGAACCAAGTAGACTTCCAAGTAAGCAATCTAAAAATTACTAGATTA
TTTAACAAGCTGCGATTCAAAATACTTGAAC

SEQ.I.D. No. 24

Intron 10 antisense sequence (172 bp).

CTGCAAAGTGAAGTAACTAATCAGTACAGCTATTACCGAATTTGACCAGCTATTGGATTAAATAATATG
AAATCCATCAAGAAATGGAAGGTAAAAAGGTTTCTACTTGTCCTTGGATAGAATTAAAGCACTTCA
TAAACCCAACACTTTCAACTTTAGATGATTTTAC

SEQ.I.D. No. 25

Intron 11 antisense sequence (145 bp).

CTGTTTTCGTCATGCGAGGATCAGAAAAAGAGTTAAATTAGACAATGTGAAAATGATTTGTTTCAGTT ACTTCTCCATAAAACTTGTTCAGTACATTAAAAACAAGCAGGAGCAATAATTTCATGGATAAGTAAAACA TATATAC

SEQ.I.D. No. 26

Intron 12 antisense sequence (242 bp).

CTGTCATGAGAACAGATTGTATGTCAGCATGAAGACAAAGATCATCAATAAACAGTTTTCTCCTTTTTG
AATTAGCTAAACAACGCAGGGGGAGGGCAGGAGGCTCAAACACTTCCGAACTCAGACAGTCGGATATCT
TATACAACTAAAGATGGATGAGACAATTACAGTTCTTTTTTGGTGAGAGAACTGTACCCTACATCTGTTA
TCTTATTATCAAAAGTTATTCAAGCAAATCCTTAC

SEQ.I.D. No. 27

Intron 13 antisense sequence (797 bp).

SEQ.I.D. No. 28
Antisense DNA sequence of the SBE gene promoter region.

CGAATGGGTTTTGATAAAACTTTGAAATTAATTTCCATTGATTAAATTATGGTACTTTGC	60
TTCAGCTGCTGCCTTCTTGTATGTTCTGATTCTTGATTTCCTCATTTTAGTGGCTTTTTA	120
TAAAAAAACATTATGACCCTTTTGTTAGTCCTCCCCTTTCTGAATATTTCACTCAGACCC	180
${\tt CATTAGTTTCGAAATTGAAGTAAAACATATTTTTTTTTT$	240
${\tt CTACTTACTTCGTTATACAATTTCTATTTAGGTAATCTAATCGACTTTTTGTATACA}$	300
TAATACATGTATTTTTGGTAAAGAGTTTTTTACTTTCTCCTAGTGGTAAGGCAGATATAG	360
${\tt TTAAGGATTTATTGACCTAATATGAACGCCAATAATTTTATATTTTGTATATACGTATAT}$	420
${\tt TTAAAAGTTTACTAGATATGTATAAATAAGATATTTAAAATTTAAATACAAAT{\tt G}$	480
${\tt ATTATGGTAAAATTTTGACCTCCAAATTAAAATATTTAAAATCAAGATTTGTCACTACTT}$	540
ATATATATCTTGTTGTAAATCCCTTTTAATCAAGTTGTGAGTTTACAAATATTCGTTGGT	600
TAGGCTAAAAAAATAAGCTATAAAGATCAAGTATAAAATTATGCATTTTCTGCATTTAA	660
TTTGGAAAAATATGTTGGAGCAATCTAAAATTGTTTGTTGATTTATAAATAA	720
TTGTTTTTAATAATTGATAAACTATTTATTCTGCTTAAAGTTTTAGAATGTCAAAAAATA	780
ATTTATTTTAATGACCTTAAATGATTGAATAAGATGTAGACACACTCAATTACAAAGTTA	840
CAATATTAATACACTTGTCTATTGGGTCATGGATTATATCATCTAATATAAATAA	900
CAAATTAAAGCTTCTTATAAAGTTCATAGGAACTAAGATAAACTTTGTGAATGGCCAAGC	960
ATTTTTCAGAACATCATGGGTGGTATGACAATCAAATTGAACTTATGGGATGAAAAATGA	1020
${\tt ATATCATTCAACTAAGAGGGCACAACTTGACATGTTAGAAAGTAAAGCAAATTTAGTAGT}$	1080
GGGCCAAATAAAAGAAATTAATTTGTCAGTTTATTCTTAAACTTTACCTTCTTTGAACTT	1140
${\tt CCACGTTATCAAAGGTTCACGGTTCATATGAAGGCCATGTGTATCCTTTTTAATTTTGGT}$	1200
ATTCCGTGTTCAATATCGATTAATTTAAATTCGCATGACAAAATCCTATATTAAAGTATA	1260
AAGTATTTTCTAAAACAGACAAGTTCAATACTTTAATTTTACACTGAATGCATAAATTTA	1320
CACTATAATAATTCCAGTCGCAGTCTACATTACAATAATTAACAATTTTAGCATGAAATG	1380
AAAAACTTTAAATTATATGCCATCAAATCACTTAAAGTATACATTTTTTTAATAACTAGT	1440
TCTAATCCCACTTGAAATGAGAGTTATTTTAATATCGACCGTTAATTACCATTTTATTAT	1500
TAAATCTGCAACTACAGTCAACTACACCAATGATTTTGCTGATGCCAACTCATAATATAA	1560
TATCCACCGTTCATGTGATTAATTCAATATTTCATATACGTACG	1620
ATTAACGTTGGATATACCATACCCTAAGCTCTGCCAAATGTCAATGTTCTATCATTAGCT	1680
ATTTTTATGCATCTATAATAGATGTTAAATTCATATTCTAAGATTGAACTTAATCATAAA	1740
CTCAAAATTTGTGGTACCTGTCAATGCCTCCAAAAGTTGATTGA	1800
CTGTGTACTTGTCTTTTCCTTGTAATAATGTATGTATGATAATAATAATAAGAGAACAAA	1860
ATATGGCAAAATAAACACTTTTTTAACATGTAACTCAAAACAAGTAATAGGCAAAAGTAC	1920
AGATGACAACACACTGTAAACATCATTGAGGAAAACCAAAAACCATACAACATTTTGA	1980
CTGTAAATGAAGAGTTTGAAAACAAAAACTATGTTCAAACCGACGCCAAGCTAACGAAAA	2040
TAGCCATAGAGTTCTAAGAAGCAGATGCAACAGTTCCACGGGTTAGTATCGTCTGTAGTA	2100
GGACCGGTCATGAGAACTCGAAAGAATCTGAAAGGAAGTAATGCATTTGAACCAGTAATT	2160
GGCCATGAT	

SEQ.I.D. No. 29 Genomic SBE gene

	*					
ATCATGGCCA	ATTACTGGTT	CAAATGCATT	ACTTCCTTTC	AGATTCTTTC	GAGTTCTCAT	60
GACCGGTCCT	ACTACAGACG	ATACTAACCC	GTGGAACTGT	TGCATCTGCT	TCTTAGAACT	120
CTATGGCTAT	TTTCGTTAGC	TTGGCGTCGG	TTTGAACATA	GTTTTTGTTT	TCAAACTCTT	180
CATTTACAGT	CAAAATGTTG	TATGGTTTTT	GTTTTCCTCA	ATGATGTTTA	CAGTGTTGTG	240
TTGTCATCTG	TACTTTTGCC	TATTACTTGT	TTTGAGTTAC	ATGTTAAAAA	AGTGTTTATT	300
TTGCCATATT	TTGTTCTCTT	ATTATTATTA	TCATACATAC	ATTATTACAA	GGAAAAGACA	360
AGTACACAGA	TCTTAACGTT	TATGTTCAAT	CAACTTTTGG	AGGCATTGAC	AGGTACCACA	420
AATTTTGAGT	TTATGATTAA	GTTCAATCTT	AGAATATGAA	TTTAACATCT	ATTATAGATG	480
CATAAAAATA	GCTAATGATA	GAACATTGAC	ATTTGGCAGA	GCTTAGGGTA	TGGTATATCC	540
AACGTTAATT	TAGTAATTTT	TGTTACGTAC	GTATATGAAA	TATTGAATTA	ATCACATGAA	600
CGGTGGATAT	TATATTATGA	GTTGGCATCA	GCAAAATCAT	TGGTGTAGTT	GACTGTAGTT	660
GCAGATTTAA	TAATAAAATG	GTAATTAACG	GTCGATATTA	AAATAACTCT	CATTTCAAGT	720
GGGATTAGAA	CTAGTTATTA	AAAAAATGTA	TACTTTAAGT	GATTTGATGG	CATATAATTT	780
AAAGTTTTTC	ATTTCATGCT	AAAATTGTTA	ATTATTGTAA	TGTAGACTGC	GACTGGAATT	840
ATTATAGTGT	AAATTTATGC	ATTCAGTGTA	AAATTAAAGT	ATTGAACTTG	TCTGTTTTAG	900
AAAATACTTT	ATACTTTAAT	ATAGGATTTT	GTCATGCGAA	TTTAAATTAA	TCGATATTGA	960
ACACGGAATA	CCAAAATTAA	AAAGGATACA	CATGGCCTTC	ATATGAACCG	TGAACCTTTG	1020
ATAACGTGGA	AGTTCAAAGA	AGGTAAAGTT	TAAGAATAAA	CTGACAAATT	AATTTCTTTT	1080
ATTTGGCCCA	CTACTAAATT	TGCTTTACTT	TCTAACATGT	CAAGTTGTGC	CCTCTTAGTT	1140
GAATGATATT	CATTTTTCAT	CCCATAAGTT	CAATTTGATT	GTCATACCAC	CCATGATGTT	1200
CTGAAAAATG	CTTGGCCATT	CACAAAGTTT	ATCTTAGTTC	CTATGAACTT	TATAAGAAGC	1260
TTTAATTTGA	CATGTTATTT	ATATTAGATG	ATATAATCCA	TGACCCAATA	GACAAGTGTA	1320
TTAATATTGT	AACTTTGTAA	TTGAGTGTGT	CTACATCTTA	TTCAATCATT	TAAGGTCATT	1380
TTAAATAAAA	ATTTTTTGAC	ATTCTAAAAC	TTTAAGCAGA	ATAAATAGTT	TATCAATTAT	1440
TAAAAACAAA	AAACGACTTA	TTTATAAATC	AACAAACAAT	TTTAGATTGC	TCCAACATAT	1500
TTTTCCAAAT	TAAATGCAGA	AAATGCATAA	TTTTATACTT	GATCTTTATA	GCTTATTTTT	1560
TTTAGCCTAA	CCAACGAATA	TTTGTAAACT	CACAACTTGA	TTAAAAGGGA	TTTACAACAA	1620
GATATATATA	AGTAGTGACA	AATCTTGATT	TTAAAATT	TAATTTGGAG	GTCAAAATTT	1680
TACCATAATC	ATTTGTATTT	TAAATTAAAT	TTTAAATATC	TTATTTATAC	ATATCTAGTA	1740
AACTTTTAAA	TATACGTATA	TACAAAATAT	AAAATTATTG	GCGTTCATAT	TAGGTCAATA	1800
AATCCTTAAC	TATATCTGCC	TTACCACTAG	GAGAAAGTAA	AAAACTCTTT	ACCAAAAATA	1860
CATGTATTAT	GTATACAAAA	AGTCGATTAG	ATTACCTAAA	TAGAAATTGT	ATAACGAGTA	1920
AGTAAGTAGA	AAAAATATAA	AACTACAATA	СТААААААА	TATGTTTTAC	TTCAATTTCG	1980
AAACTAATGG	GGTCTGAGTG	AAATATTCAG	AAAGGGGAGG	ACTAACAAAA	GGGTCATAAT	2040
GTTTTTTTAT	AAAAAGCCAC	TAAAATGAGG	AAATCAAGAA	TCAGAACATA	CAAGAAGGCA	2100
GCAGCTGAAG	CAAAGTACCA	TAATTTAATC	AATGGAAATT	AATTTCAAAG	TTTTATCAAA	2160
ACCCATTCGA	GGATCTTTTC	CATCTTTCTC	ACCTAAAGTT	TCTTCAGGGG	TAATTTTTAC	2220
TAATTTCATG	TTAATTTCAA	TTATTTTTAG	CCTTTGCATT	TCATTTTCCA	ATATATCTGG	2280
ATCATCTCCT	TAGTTTTTTA	TTTTATTTT	TATAATATCA	AATATGGAAG	AAAAATGACA	2340
CTTGTAGAGC	CATATGTAAG	TATCATGTGA	CAAATTTGCA	AGGTGGTTGA	GTGTATAAAA	2400
TTCAAAAATT	GAGAGATGGA	GGGGGGTGG	GGGBARAGAC	AATATTTAGA	AAGAGTGTTC	2460
TAGGAGGTTA	TGGAGGACAC	GGATGAGGGG	TAGAAGGTTA	GTTAGGTATT	TGAGTGTTGT	2520

	•					
CTGGCTTATC	CTTTCATACT	AGTAGTCGTG	GAATTATTTG	GGTAGTTTCT	TGTTTTGTTA	2580
TTTGATCTTT	GTTATTCTAT	TTTCTGTTTC	TTGTACTTCG	ATTATTGTAT	TATATATCTT	2640
GTCGTAGTTA	TTGTTCCTCG	GTAAGAATGC	TCTAGCATGC	TTCCTTTAGT	GTTTTATCAT	2700
GCCTTCTTTA	TATTCGCGTT	GCTTTGAAAT	GCTTTTACTT	TAGCCGAGGG	TCTATTAGAA	2760
ACAATCTCTC	TATCTCGTAA	GGTAGGGGTA	AAGTCCTCAC	CACACTCCAC	TTGTGGGATT	2820
ACATTGTGTT	TGTTGTTGTA	AATCAATTAT	GTATACATAA	TAAGTGGATT	TTTTACAACA	2880
CAAATACATG	GTCAAGGGCA	AAGTTCTGAA	CACATAAAGG	GTTCATTATA	TGTCCAGGGA	2940
TATGATAAAA	ATTGTTTCTT	TGTGAAAGTT	ATATAAGATT	TGTTATGGCT	TTTGCTGGAA	3000
ACATAATAAG	TTATAATGCT	GAGATAGCTA	CTGAAGTTTG	TTTTTTCTAG	CCTTTTAAAT	3060
GTACCAATAA	TAGATTCCGT	ATCGAACGAG	TATGTTTTGA	TTACCTGGTC	ATGATGTTTC	3120
TATTTTTAC	ATTTTTTGG	TGTTGAACTG	CAATTGAAAA	TGTTGTATCC	TATGAGACGG	3180
ATAGTTGAGA	ATGTGTTCTT	TGTATGGACC	TTGAGAAGCT	CAAACGCTAC	TCCAATAATT	3240
TCTATGAATT	CAAATTCAGT	TTATGGCTAC	CAGTCAGTCC	AGAAATTAGG	ATATGCTGCA	3300
TATACTTGTT	CAATTATACT	GTAAAATTTC	TTAAGTTCTC	AAGATATCCA	TGTAACCTCG	3360
		CTAGAAATAA				3420
		AACGGTCTTG			•	3480
•		GTTTGATAAT				3540
		GTGCTCATGG				3600
		TCCGCCAATT				3660
TTTACTCCTG	TGGGCATTTG	TTATTTGAAT	TACAATCTTT	ATGAGCATGG	TGTTTTCACA	3720
TTATCAACTT	CTTTCATGTG	GTATATAACA	GTTTTTAGCT	CCGTTAATAC	CTTTCTTCTT	3780
TTTGATATAA	ACTAACTGTG	GTGCATTGCT	TGCBKKKATG	AAGCACAGTT	CAGCTATTTC	3840
CGCTGTTTTG	ACCGATGACG	ACAATTCGAC	AATGGCACCC	CTAGAGGAAG	ATGTCAAGAC	3900
TGAAAATATT	GGCCTCCTAA	ATTTGGATCC	AACTTTGGAA	CCTTATCTAG	ATCACTTCAG	3960
ACACAGAATG	AAGAGATATG	TGGATCAGAA	AATGCTCATT	GAAAAATATG	AGGGACCCCT	4020
TGAGGAATTT	GCTCAAGGTA	ACAGCCAAAA	GTTGTGCTTT	AGGCAGTTTG	ACCTTATTTT	4080
GGAAGATGAA	TTGTTTATAC	CTACTTTGAC	TTTGCTAGAG	AATTTTGCAT	ACCGGGGAGT	4140
AAGTAGTGGC	TCCATTTAGG	TGGCACCTGG	CCATTTTTTT	GATCTTTTAA	AAAGCTGTTT	4200
GATTGGGTCT	TCAAAAAAGT	AGACAAGGTT	TTTGGAGAAG	TGACACACCC	CCGGAGTGTC	4260
AGTGGCAAAG	CAAAGATTTT	CACTAAGGAG.	ATTCAAAATA	TAAAAAAAGT	ATAGACATAA	4320
AGAAGCTGAG	GGGATTCAAC	ATGTACTATA	CAAGCATCAA	ATATAGTCTT	AAAGCAATTT	4380
TGTAGAAATA	AAGAAAGTCT	TCCTTCTGTT	GCTTCACAAT	TTCCTTCTAT	TATCATGAGT	4440
TACTCTTTCT	GTTCGAAATA	GCTTCCTTAA	TATTAAATTC	ATGATACTTT	TGTTGAGATT	4500
TAGCAGTTTT	TTCTTGTGTA	AACTGCTCTC	TTTTTTTGCA	GGTTATTTAA	AATTTGGATT	4560
CAACAGGGAA	GATGGTTGCA	TAGTCTATCG	TGAATGGGCT	CCTGCTGCTC	AGTAGGTCCT	4620
CGTCTACTAC	AAAATAGTAG	TTTCCATCAT	CATAACAGAT	TTTCCTATTA	AAGCATGATG	4680
TTGCAGCATC	ATTGGCTTTC	TTACATGTTC	TAATTGCTAT	TAAGGTTATG	CTTCTAATTA	4740
ACTCATCCAC	AATGCAGGGA	AGCAGAAGTT	ATTGGCGATT	TCAATGGATG	GAACGGTTCT	4800
AACCACATGA	TGGAGAAGGA	CCAGTTTGGT	GTTTGGAGTA	TTAGAATTCC	TGATGTTGAC	4860
AGTAAGCCAG	TCATTCCACA	CAACTCCAGA	GTTAAGTTTC	GTTTCAAACA	TGGTAATGGA	4920
	•				CACAAAGTTT	4980
					TTGTTATTCA	5040
		*			GTTCTTACTA	5100
					TTTACTAATT	5160
TGGACTTGAT	CCCATACTCT	TTCCCTTAAC	AAAATGAGTC	AATTCTATAA	GTGCTTGAGA	5220

	•					
ACTTACTACT	TCAGCAATTA	AACAGGTACC	ACTTCAAATA	CCCTCGCCCT	CCCAAACCCC	5280
GAGCCCCACG	AATCTATGAA	GCACATGTCG	GCATGAGCAG	CTCTGAGCCA	CGTGTAAATT	5340
CGTATCGTGA	GTTTGCAGAT	GATGTTTTAC	CTCGGATTAA	GGCAAATAAC	TATAATACTG	5400
TCCAGTTGAT	GGCCATAATG	GAACATTCTT	ACTATGGATC	ATTTGGATAT	CATGTTACAA	5460
ACTTTTTTGC	TGTGAGCAGT	AGATATGGAA	ACCCGGAGGA	CCTAAAGTAT	CTGATAGATA	5520
AAGCACATAG	CTTGGGTTTA	CAGGTTCTGG	TGGATGTAGT	TCACAGTCAT	GCAAGCAATA	5580
ATGTCACTGA	TGGCCTCAAT	GGCTTTGATA	TTGGCCAAGG	TTCTCAAGAA	TCCTACTTTC	5640
ATGCTGGAGA	GCGAGGGTAC	CATAAGTTGT	GGGATAGCAG	GCTGTTCAAC	TATGCCAATT	5700
GGGAGGTTCT	TCGTTTCCTT	CTTTCCAACT	TGAGGTGGTG	GCTAGAAGAG	TATAACTTTG	5760
ACGGATTTCG	ATTTGATGGA	ATAACTTCTA	TGCTGTATGT	TCATCATGGA	ATCAATATGG	5820
GATTTACAGG	AAACTATAAT	GAGTATTTCA	GCGAGGCTAC	AGATGTTGAT	GCTGTGGTCT	5880
ATTTAATGTT	GGCCAATAAT	CTGATTCACA	AGATTTTCCC	AGATGCAACT	GTTATTGCCG	5940
AAGATGTTTC	TGGTATGCCG	GGCCTTGGCC	GGCCTGTTTC	TGAGGGAGGA	ATTGGTTTTG	6000
TTTACCGCCT	GGCAATGGCA	ATCCCAGATA	AGTGGATAGA	TTATTTAAAG	AATAAGAATG	6060
ATGAAGATTG	GTCCATGAAG	GAAGTAACAT	CGAGTTTGAC	AAATAGGAGA	TATACAGAGA	6120
AGTGTATAGC	ATATGCGGAG	ACCCATGATC	AGGTATTTTA	AATTTATTTC	TACAACTAAA	6180
TAATTCTCAG	AACAATTGTT	AGATAGAATC	CAAATATATA	CGTCCTGAAA	GTATAAAAGT	6240
ACTTATTTTC	GCCATGGGCC	TTCAGAATAT	TGGTAGCCGC	TGAATATCAT	GATAAGTTAT	6300
TTATCCAGTG	ACATTTTAT	GTTCACTCCT	ATTATGTCTG	CTGGATACAG	TCTATTGTTG	6360
GTGACAAGAC	CATTGCATTT	CTCCTAATGG	ACAAAGAGAT	GTATTCTGGC	ATGTCTTGCT	6420
TGACAGATGC	TTCTCCTGTT	GTTGATCGAG	GAATTGCGCT	TCACAAGGTT	TGTCTGTTTC	6480
TATTGCATTT	TAAGGTTCAT	ATAGGTTAGC	CACGGAAAAT	CTCACTCTTT	GTGAGGTAAC	6540
CAGGGTTCTG	ATGGATTATT	CAATTTTCTC	GTTTATCATT	TGTTTATTCT	TTTCATGCAT	6600
TGTGTTTCTT	TTTCAATATC	CCTCTTATTT	GGAGGTAATT	TTTCTCATCT	ATTCACTTTT	6660
AGCTTCTAAC	CACAGATGAT	CCATTTTTTC	ACAATGGCCT	TGGGAGGAGA	GGGGTACCTC	6720
AATTTCATGG	GTAACGAGGT	ATGTCTTACA	TCTTTAGATA	TTTTGTGATA	ATTACAATTA	6780
GTTTGGCTTA	CTTGAACAAG	ATTCATTCCT	CAAAATGACC	TGAACTGTTG	AACATCAAAG	6840
GGGTTGAAAC	ATAGAGGAAA	ACAACATGAT	GAATGTTTCC	ATTGTCTAGG	GATTTCTATT	6900
ATGTTGCTGA	GAACAAATGT	CATCTTAAAA	AAAACATTGT	TTACTTTTTT	GTAGTATAGA	6960
AGATTACTGT	ATAGAGTTTG	CAAGTGTGTC	TGTTTTGGAG	TAATTGTGAA	ATGTTTGATG	7020
					AATAATTGGA	7080
					AGATACAAGG	7140
				•	TGCTTACTTG	
					ATTTTCCAAC	7260
	•				TATTACCGCC	
•					GCTATAGTTT	
					TTACAATGAA	
					TCCTCCCCT	
			•		CGATGAAAAG	
					TAATAAGGTA	7620
					CCAAGGACAA	•
					AATCCAATAG	
			+		TTGTTGTGTT	
					ACGAAGGGTA	
TATATGTTTT	ACTTATCCAT	GAAATTATTG	CTCTGCTTGT	TTTTAATGTA	CTGAACAAGT	7920

TTTATGGAGA	AGTAACTGAA	ACAAATCATT	TTCACATTGT	CTAATTTAAC	TCTTTTTTCT	7980
GATCCTCGCA	TGACGAAAAC	AGGTATAAAG	TTGGATGTGA	CTTGCCAGGG	AAGTACAGAG	8040
TTGCACTGGA	CAGTGATGCT	TGGGAATTTG	GTGGCCATGG	AAGAGTAAGG	ATTTGCTTGA	8100
ATAACTTTTG	ATAATAAGAT	AACAGATGTA	GGGTACAGTT	CTCTCACCAA	AAAGAACTGT	8160
AATTGTCTCA	TCCATCTTTA	GTTGTATAAG	ATATCCGACT	GTCTGAGTTC	GGAAGTGTTT	8220
GAGCCTCCTG	CCCTCCCCCT	GCGTTGTTTA	GCTAATTCAA	AAAGGAGAAA	ACTGTTTATT	8280
GATGATCTTT	GTCTTCATGC	TGACATACAA	TCTGTTCTCA	TGACAGACTG	GTCATGATGT	8340
TGACCATTTC	ACATCACCAG	AAGGAATACC	TGGAGTTCCA	GAAACAAATT	TCAATGGTCG	8400
TCCAAATTCC	TTCAAAGTGC	TGTCTCCTGC	GCGAACATGT	GTGGTACAGT	TCTTGCCGTG	8460
TGACCTCCCT	TTTTATTGTG	GTTTTGTTCA	TAGTTATTTG	AATGCGATAG	AAGTTAACTA	8520
TTGATTACCG	CCACAATCGC	CAGTTAAGTC	CTCTGAACTA	CTAATTTGAA	AGGTAGGAAT	8580
AGCCGTAATA	AGGTCTACTT	TTGGCATCTT	ACTGTTACAA	AACAAAAGGA	TGCCAAAAAA	8640
ATTCTTCTCT	ATCCTCTTTT	TCCCTAAACC	AGTGCATGTA	GCTTGCACCT	GCATAAACTT	8700
AGGTAAATGA	TCAAAAATGA	AGTTGATGGG	AACTTAAAAC	CGCCCTGAAG	TAAAGCTAGG	8760
AATAGTCATA	TAATGTCCAC	CTTTGGTGTC	TGCGCTAACA	TCAACAACAA	CATACCTCGT	8820
GTAGTCCCAC	AAAGTGGTTT	CAGGGGGAGG	GTAGAGTGTA	TGCAAAACTT	ACTCCTATCT	8880
CAGAGGTAGA	GAGGATTTTT	TCAATAGACC	CTTGGCTCAA	GAAAAAAGT	CCAAAAAGAA	8940
GTAACAGAAG	TGAAAGCAAC	ATGTGTAGCT	AAAGCGACCC	AACTTGTTTG	GGACTGAAGT	9000
AGTTGTTGTT	GTTGAAACAG	TGCATGTAGA	TGAACACATG	TCAGAAAATG	GACAACACAG	9060
TTATTTTGTG	CAAGTCAAAA	AAATGTACTA	CTATTTCTTT	GTGCAGCTTT	ATGTATAGAA	9120
AAGTTAAATA	ACTAATGAAT	TTTGCTAGCA	GAAAAATAGC	TTGGAGAGAA	ATTTTTTTA	9180
TTGAACTAAG	CTAACTATAT	TCATCTTTCT	TTTTGCTTCT	TCTTCTCCTT	GTTTGTGAAG	9240
GCTTATTACA	GAGTTGATGA	ACGCATGTCA	GAAACTGAAG	ATTACCAGAC	AGACATTTGT	9300
AGTGAGCTAC	TACCAACAGC	CAATATCGAG	GAGAGTGACG	AGAAACTTAA	AGATTCGTTA	9360
TCTACAAATA	TCAGTAACAT	TGACGAACGC	ATGTCAGAAA	CTGAAGTTTA	CCAGACAGAC	9420
ATTTCTAGTG	AGCTACTACC	AACAGCCAAT	ATTGAGGAGA	GTGACGAGAA	ACTTAAAGAT	9480
TCGTTATCTA	CAAATATCAG	TAACATTGAT	CAGACTGTTG	TAGTTTCTGT	TGAGGAGAGA	9540
GACAAGGAAC	TTAAAGATTC	ACCGTCTGTA	AGCATCATTA	GTGATGTTGT	TCCAGCTGAA	9600
TGGGATGATT	CAGATGCAAA	CGTCTGGGGT	GAGGACTAGT	CAGATGATTG	ATCGACCCTT	9660
CTACCGATTG	GTGATCGCTA	TCCTTGCTCT	CTGAGAAATA	GGTGAGGCGA	AACAAAAAAT	9720
AATTTGCATG	ATAAAAAGTC	TGATTTTATG	ATCGCTATCC	TCGCTCTCTG	AGAAAGAAGC	9780
					CCTGGGACTC	9840
					TTAGTTAAGA	9900
				,	GATAATTATA	9960
					TCTATCAAAA	10020
					TAATTACATC	10080
					TTTGTTGCCA	10140
					ATTTAGAGAT	10200
					CATGCATGAA	10260
					ATCATTAATG	10320
					GTAAAAATAA	10380
					TAAATCTTAC	10440
					A CGCGTCCTGA	
					TTAGGTATGC	10560
CAAAAAAAA	AACACGATCO	TTTGCACCC	F TTCGATGAT	r atcagtatg:	r tcacaaaaaa	10620

	AACTTAAGTT	CATCCCAGTG	TACAACAGCC	CCAACATCTG	CCCCAAGTAA	CAAAAAACAA	10680
	CCAATTTATC	TTATTCTTAT	CTGCCACAAA	ATAATCGGTT	TCACACTATT	CTCTTGTTAT	10740
	ACAAAATTGA	CAAGTAGGAA	GGAGAGGAGT	CATCCAAATA	AACGGTGCAC	GTTCTTTGAG	10800
	AAAAGTCTTA	TTTTTCGTAA	GATCCAATTT	CAACAAACTT	TTCTTCAAGT	CAAAATTCCT	10860
1	GATAGTGTAT	CTCCTCTCGA	CGACCTCTTG	CATTGAACGA	TCTCCGCTTA	TCATGAAAAG	10920
•	TTGCTTGGAT	AACAAGTATT	GCAAGGGGG	GACAGTAGCT	ATTAAGTTAG	TCGGCCCAAG	10980
(GAAATGGAGG	AGTGATAGTC	TCGAATATTA	TTCACCTCTT	TAGCATTACC	CGGTCTGGCT	11040
•	TTAAGGAGTT	ACGTCTTTTA	CGCTCGCCAA	TTTCTTTTTT	TAGAATGGTT		11100
	CGCGAGTTG	TGGAAGGTTC	AAGTTACTCG	ATTCGTGATT	TTCAAGTATG	AGTGGTGAGA	11160
(GAGATTCGAT	ATTTTCACGA	GGTGTATTCG	AGGTCTAGTA	GAACGAAGGG	TGTCACTAAT	11220
(SAAAGTTTCA	AGAGTTCATC	ATCATCTTCT	TCTAGTAGAT	TTTCGCTTTC	AAATGAGTAT	11280
(SAAAATTCTT	CCTCTTTTCT	ATTGATTTTC	TTCATTGTTT	TCTTCATTGT	TGTGGTTGTT	11340
7	ATTGAAAAGA	AAGAAAATTT	ATAACAGAAA	AAGATGTCAA	AAAAAAGGTA	AAATGAAAGA	11400
C	TATCATATA	CTTAAAGAGT	TGCGTAGAGA	TAAGTCAAAA	GAAACAGAAT	TATAGTAATT	11460
	CAGCTAAGT						11478

5

25

CLAIMS

- 1. A method of affecting enzymatic activity in a plant (or a cell, a tissue or an organ thereof) comprising expressing in the plant (or a cell, a tissue or an organ thereof) a nucleotide sequence wherein the nucleotide sequence codes, partially or completely, for an intron in a sense orientation; and wherein the nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron.
- 2. A method according to claim 1 wherein starch branching enzyme activity is affected and/or wherein the levels of amylopectin are affected and/or the composition of starch is changed.
- 3. A method of affecting enzymatic activity in a starch producing organism (or a cell, a tissue or an organ thereof) comprising expressing in the starch producing organism (or a cell, a tissue or an organ thereof) a nucleotide sequence wherein the nucleotide sequence codes, partially or completely, for an intron in a sense orientation; wherein the nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron; and wherein starch branching enzyme activity is affected and/or the levels of amylopectin are affected and/or the composition of starch is changed.
 - 4. A method according to any one of claims 1 to 3 wherein the nucleotide sequence does not contain a sequence that is sense to an exon sequence.
 - 5. A method according to any one of the preceding claims wherein the enzymatic activity is reduced or eliminated.
- 6. A method according to any one of the preceding claims wherein the nucleotide sequence codes for at least substantially all of at least one intron in a sense orientation.

10


20

30

- 7. A method according to any one of the preceding claims wherein the nucleotide sequence codes for all of at least one intron in a sense orientation.
- 8. A method according to any one of the preceding claims wherein the nucleotide sequence comprises the sequence shown as any one of SEQ.I.D. No. 1 to SEQ.I.D. No. 13 or a variant, derivative or homologue thereof, including combinations thereof.
 - 9. A method according to any one of the preceding claims wherein the nucleotide sequence is expressed by a promoter having a sequence shown as SEQ.I.D. No. 14 or a variant, derivative or homologue thereof.
 - 10. A sense sequence comprising the nucleotide sequence as defined in claim 8 or a variant, derivative or homologue thereof.
- 15 11. A promoter having a sequence shown as SEQ.I.D. No. 14, or a variant, derivative or homologue thereof.
 - 12. A promoter according to claim 11 in combination with a gene of interest ("GOI").
 - 13. A construct capable of comprising or expressing the invention according to any one of claims 10 to 12.
- 14. A vector comprising or expressing the invention according to any one of claims 10 to 13.
 - 15. A combination of nucleotide sequences comprising a first nucleotide sequence coding for a recombinant enzyme; and a second nucleotide sequence which corresponds to an intron in a sense orientation; wherein the intron is an intron that is associated with a genomic gene encoding an enzyme corresponding to the recombinant enzyme; and wherein the second nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron.

5

- 16. A cell, tissue or organ comprising or expressing the invention according to any one of claims 10 to 15.
- 17. A transgenic starch producing organism comprising or expressing the invention according to any one of claims 10 to 16.
 - 18. A transgenic starch producing organism according to claim 17 wherein the organism is a plant.
- 19. A starch obtained from the invention according to any one of the preceding claims.
 - 20. pBEA11 (NCIMB 40754).
- 15 21. An intron nucleotide sequence that is obtainable from λ -SBE 3.2 (NCIMB 40751) or λ -SBE 3.4 (NCIMB 40752) or a variant, derivative or homologue thereof.
- 22. A method of expressing a recombinant protein or enzyme in a host organism comprising expressing a nucleotide sequence coding for the recombinant protein or enzyme; and expressing a further nucleotide sequence; wherein the further nucleotide sequence codes, partially or completely, for an intron in a sense orientation; wherein the intron is an intron normally associated with the genomic gene encoding a protein or an enzyme corresponding to the recombinant protein or enzyme; and wherein the further nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron.






Fig 1




Fig 2




Fig 3




Fig 4




Fig 5




Fig 6




Fig 7

	10 20 30 40 50 60 12345678901234567890123456789012345678901234567890
60	ATCATGGCCAATTACTGGTTCAAATGCATTACTTCCTTTCAGATTCTTTCGAGTTCTCAT
120	GACCGGTCCTACTACAGACGATACTAACCCGTGGAACTGTTGCATCTGCTTCTTAGAACT
180	CTATGGCTATTTTCGTTAGCTTGGCGTCGGTTTGAACATAGTTTTTGTTTTCAAACTCTT
240	CATTTACAGTCAAAATGTTGTATGGTTTTTGTTTTCCTCAATGATGTTTACAGTGTTGTG
300	TTGTCATCTGTACTTTTGCCTATTACTTGTTTTGAGTTACATGTTAAAAAAAGTGTTTATT
360	TTGCCATATTTTGTTCTCTTATTATTATTATCATACATAC
420	AGTACACAGATCTTAACGTTTATGTTCAATCAACTTTTGGAGGCATTGACAGGTACCACA
480	AATTTTGAGTTTATGATTAAGTTCAATCTTAGAATATGAATTTAACATCTATTATAGATG
540	CATAAAAATAGCTAATGATAGAACATTGACATTTGGCAGAGCTTAGGGTATGGTATATCC
600	AACGTTAATTTAGTAATTTTTGTTACGTACGTATATGAAATATTGAATTAATCACATGAA
660	CGGTGGATATTATATTATGAGTTGGCATCAGCAAAATCATTGGTGTAGTTGACTGTAGTT
720	GCAGATTTAATAATAAATGGTAATTAACGGTCGATATTAAAATAACTCTCATTTCAAGT
780	GGGATTAGAACTAGTTATTAAAAAAATGTATACTTTAAGTGATTTGATGGCATATAATTT
840	AAAGTTTTTCATTTCATGCTAAAATTGTTAATTATTGTAATGTAGACTGCGACTGGAATT
900	ATTATAGTGTAAATTTATGCATTCAGTGTAAAATTAAAGTATTGAACTTGTCTGTTTTAG
960	AAAATACTTTATACTTTAATATAGGATTTTGTCATGCGAATTTAAATTAATCGATATTGA
1020	ACACGGAATACCAAAATTAAAAAGGATACACATGGCCTTCATATGAACCGTGAACCTTTG
1080	ATAACGTGGAAGTTCAAAGAAGGTAAAGTTTAAGAATAAACTGACAAATTAATT
1140	ATTTGGCCCACTACTAAATTTGCTTTACTTTCTAACATGTCAAGTTGTGCCCTCTTAGTT
1200	GAATGATATTCATTTTCATCCCATAAGTTCAATTTGATTGTCATACCACCCATGATGTT
1260	CTGAAAAATGCTTGGCCATTCACAAAGTTTATCTTAGTTCCTATGAACTTTATAAGAAGC
1320	ITTAATTTGACATGTTATTTATATTAGATGATATAATCCATGACCCAATAGACAAGTGTA
1380	TTAATATTGTAACTTTGTAATTGAGTGTGTCTACATCTTATTCAATCATTTAAGGTCATT
1440	AAAATAAATTATTTTTTGACATTCTAAAACTTTAAGCAGAATAAATA
1500	FAAAAACAAAAAACGACTTATTTATAAATCAACAAACAATTTTAGATTCCTCCAACATTA

10 20 30 40 50 60 12345678901234567890123456789012345678901234567890	
TTTTCCAAATTAAATGCAGAAAATGCATAATTTTATACTTGATCTTTATAGCTTATTTTT	1560
TTTAGCCTAACCAACGAATATTTGTAAACTCACAACTTGATTAAAAGGGATTTACAACAA	1620
GATATATAAGTAGTGACAAATCTTGATTTTAAATATTTTAATTTGGAGGTCAAAATTT	1680
TACCATAATCATTTGTATTTATAATTAAATTTTAAATATCTTATTTAT	1740
AACTTTTAAATATACGTATATACAAAATATAAAATTATTGGCGTTCATATTAGGTCAATA	1800
AATCCTTAACTATATCTGCCTTACCACTAGGAGAAAGTAAAAAACTCTTTACCAAAAATA	1860
CATGTATTATGTATACAAAAGTCGATTAGATTACCTAAATAGAAATTGTATAACGAGTA	1920
AGTAAGTAGAAATATAAAAAAACTACAATACTAAAAAAAA	1980
AAACTAATGGGGTCTGAGTGAAATATTCAGAAAGGGGAGGACTAACAAAAGGGTCATAAT	2040
GTTTTTTTATAAAAGCCACTAAAATGAGGAAATCAAGAATCAGAACATACAAGAAGGCA	2100
GCAGCTGAAGCAAAGTACCATAATTTAATCAATGGAAATTAATT	2160
P I R G S F P S F S F S F S F S F S F S F S F S	2220
taatttcatgttaatttcaattatttttagcctttgcatttcattttccaatatctgg	2280
atcatctccttagttttttattttttttataatatcaaatatggaagaaaaatgaca	2340
cttgtagagccatatgtaagtatcatgtgacaaatttgcaaggtggttgagtgtataaaa	2400
ttcaaaaattgagagatggaggggggggggbaragacaatatttagaaagagtgttc	2460
taggaggttatggaggacacggatgaggggtagaaggttagtta	2520
ctggcttatcctttcatactagtagtcgtggaattatttgggtagtttcttgtttgt	2580
tttgatctttgttattctattttctgtttcttgtacttcgattattgtattatatctt	2640
gtcgtagttattgttcctcggtaagaatgctctagcatgcttcctttagtgttttatcat	2700
gccttctttatattcgcgttgctttgaaatgcttttactttagccgagggtctattagaa	2760
acaatctctctatctcgtaaggtaggggtaaagtcctcaccacactccacttgtgggatt	2820
acattgtgtttgttgttgtaaatcaattatgtatacataataagtggattttttacaaca	2880
caaatacatggtcaagggcaaagttctgaacacataaagggttcattatatgtccaggga	2940
atgataaaaattgtttctttgtgaaagttatataagatttgttatggcttttgctggaa	3000

10 20 30 40 50 60 12345678901234567890123456789012345678901234567890	
acataataagttataatgctgagatagctactgaagtttgttt	3060
gtaccaataatagattccgtatcgaacgagtatgttttgattacctggtcatgatgtttc	3120
tattttttacatttttttggtgttgaactgcaattgaaaatgttgtatcctatgagacgg	3180
atagttgagaatgtgttctttgtatggaccttgagaagctcaaacgctactccaataatt	3240
tctatgaattcaaattcagtttatggctaccagtcagtccagaaattaggatatgctgca	3300
tatacttgttcaattatactgtaaaatttcttaagttctcaagatatccatgtaacctcg	3360
agaatttctttgacagGCTTCTAGAAATAAGATATGTTTTCCTTCTCAACATAGTACTGG A S R N K I C F P S Q H S T G	3420
ACTGAAGTTTGGATCTCAGGAACGGTCTTGGGATATTTCTTCCACCCCAAAATCAAGAGT L K F G S Q E R S W D I S S T P K S R V	3480
TAGAAAAGATGAAAGGgtatgtttgataatttatatggttgcatggatagtatataaata R K D E R	3540
gttggaaaacttctggactggtgctcatggcatatttgatctgtgcaccgtgtggagatg	3600
tcaaacatgtgttacttcgttccgccaatttataataccttaacttgggaaagacagctc	3660
tttactcctgtgggcatttgttatttgaattacaatctttatgagcatggtgttttcaca	3720
ttatcaacttctttcatgtggtatataacagtttttagctccgttaatacctttcttctt	3780
tttgatataaactaactgtggtgcattgcttgcbkkkATGAAGCACAGTTCAGCTATTTC	3840
CGCTGTTTTGACCGATGACGACAATTCGACAATGGCACCCCTAGAGGGAAGATGTCAAGAC	3900
TGAAAATATTGGCCTCCTAAATTTGGATCCAACTTTGGAACCTTATCTAGATCACTTCAG	3960
E N I G L L N L D P T L E P Y L D H F R ACACAGAATGAAGAGATATGTGGATCAGAAAATGCTCATTGAAAAATATGAGGGACCCCT H R M K R Y V D Q K M L I E K Y E G P L	4020
TGAGGAATTTGCTCAAGgtaacagccaaaagttgtgctttaggcagtttgaccttatttt E E F A Q G	4080
ggaagatgaattgtttatacctactttgactttgctagagaattttgcataccggggagt	4140
aagtagtggctccatttaggtggcacctggccatttttttgatcttttaaaaagctgttt	4200
gattgggtcttcaaaaaagtagacaaggtttttggagaagtgacacacccccggagtgtc	4260
agtggcaaagcaaagattttcactaaggagattcaaaatataaaaaaagtatagacataa	4320
agaagctgaggggattcaacatgtactatacaagcatcaaatatagtcttaaagcaattt	4380
tgtagaaataaagaaagtcttccttctgttgcttcacaatttccttctattatcatgagt	4440
tactctttctgttcgaaatagcttccttaatattaaattcatgatacttttgttgagatt	4500

<u>123</u>	<u>456</u>	19 789	0 012:	345	678	20 901:	234	567	30 890	123	456	789/	0	2454	570	50	~		60	
tag	cag	ttt	ttto	ctt	ata	taa	act/	70+	C+ C	+++	300	7031	<u> </u>	3430	0/8	901	234	567	890	
		ttti CCN:																		4560
		GGAZ E	~	u	_		v	7	-	m.	IAI '	70	n -	•	•	_				4620
cgt	cta	ctad	caaa	aata	agt:	agti	ttc	cat	cat	cata	aaca	agat	ttt	cct	ati	taa	agc	atg	atg	4680
ttg	cag	cato	catt	gg	ctt	tcti	taca	atgi	ttc	taat	tg	ctat	taa	aggt	tat	gc	ttc	taa	tta	4740
		ccad			1 9 G	GAA(3CAC	GAA(STT	ATTO	GCC	GATT	rrcz							4800
AAC							• •	•. 1	~		2 1	7 7	•	7 /	•			_		
			7 E		, I) (, ,	• (. I	υv	a c	7 7	re	•		•	•		_	4860
AGT.			, ,			1 E	v :	•	₹ 1	√ Tu		? 6			, ,	• .	-		_	4920
GTG V 1						CTC	CTI	rGG/	ATA	AAG1	ATC	CC.	CTC	CAC	ACC	CC	ACA	AAG		4980
GCA(GCA(CAI	ATC	ATC	GTC	TCI	[AC]	NGG(:AC(CAC	CAC	CTI	CAC	AAA) ; \Ggt	tti	r : :gt:	K tat	F tca	5040
tac				, ,	3 Y	, ,	·	V I	, ,		, ,				•					5100
gtc																				5160
tgga																				
																				5220
acti								Y	м	100	v	v	70	-	-	-	••	_	_	5280
GAGO	ייייי	ACG R	AAT	CTA	TGA	LAGC	ACA	TGI	CGC	CAI	GAG	CAG	CTC	TGA	GCC	ACC	TG:	CAA	ATT	5340
-	₽.	- 1	_	I	E.	A	Ħ.	v	72	M		_	_	-	-	_			_	
CGTA	R.	E	F	TGC	AGA	VI GA	TGT	111	ACC	TCG	GAI	TAA	'GCC	AAA	TAA	CTA	ATA	ATA	CTG	5400
•			T.		$\boldsymbol{\omega}$	u	v	1.		-	т	~	•	3.7	••			_		
TCC		7.7	n		m		-	-	•	v	C	C	-	_	10	••		_		5460
ACTI	TTI	TGC	TGT	GAG	CAG	TAG	ATA	TGC	AAA	VCCC	೧೭೨೨	CCA	CCT	מממי	Cun y	TC I	V V	T	N	
•	-		v	_		ĸ	Y	(:	N	ם		~	•	7.0	••	-	_	_		5520
AAGC	ACA	TAG	CII	GGG	LLI	'ACA	.GGT	TCI	'GGT	'GGA	TGT	'AGT	TCA	CAG	מית	TGC	`AAC	CA	ATA	5580
• • •	••			u		•	v		v	- 11	17	7.7	u	_	**	•	-			
ATG1	T	.TGA D	166	CCT	CAA	TGG	CII	TGA	TAT	TGG	CCA	AGG	TTC	TCA	AGA	ATC	CT	CT	MC	5640
	•	U	J		7.4	G		D	•	G	n	G	•	\sim	T	_		_		
ATGC	G	E	R	AGG	OIW	יררא	TAA	GT-T	GI.	iGGA	TAG	CAG	GCT	GIT	CAA	CTA	TGC	CAZ	\TT	5700
		ىلىكىلە چ	عرت م	ىلىملىك ج	-	AU√AL Σ	ωπ. Υ	C22	~W	D	S	R	L	F	N	Y	A	N	W	
GGGA E	v	L	R	F	T.	T.	110	CAA N	T - 1	UAU P	GTG	GIG	GCT	AGA	AGA	.GTA	TA	CT.	TTG	5760
ACGG	ATT	TCG	ATT	TGA	TGG	AAT	AAC	TALC.	ጥልጥ ພ	ינהריכה צר	Λυν Μ	w ™~m	шС. Г	E	E	Y	N	F	D	
•	-	*,	4	ע	u		11		M	F .	v	\ \ T	u	11	\sim	-	••		_	5820
GATI	TW/	\sim	$\sim\sim$	CIA	TAA	TIA	(TIA	,,,,,,,	$T^*\Delta C$	$CC\Delta$	ccc	ጥእሎ	707		~~~	mac				F000
	-	G	7.4	1	1/4	Ŀ	Y	F.		F	Δ	T	₽	37	•	•		• •		5880
ATTI	Trans.	011	$\sigma \sigma$	$-\infty$	IMM	TUT	GAT	TC A	CAA	CAT	للعلمك	\sim	3 C 3	\sim	330	There	ייערא ∧	νης. •	Y	5040
	7.7	-	_	TA.	IN	1.		- н	ĸ	T		ъ.	_	•	-	• •	_	_	_	5940
AAGA	TGT	TIC	TGG'	TAT	GCC	GGG	CCT	\mathtt{TGG}	CCG	CCC	ىدىك	مالس	תבים	~~~	300	220	ALCA T	ململئ وي	באנה בי	6000
D	V	S	G	M	P	G	L	G	R	P	v	S	E	G	G			· • • •	V.	0000

10 20 30 40 50 60 12345678901234567890123456789012345678901234567890	
TTTACCGCCTGGCAATGGCAATCCCAGATAAGTGGATAGATTATTTAAAGAATAAGAATG	
	6060
ATGAAGATTGGTCCATGAAGGAAGTAACATCGAGTTTGACAAATAGGAGATATACAGAGA E D W S M K E V T S S L T N R R Y T E K	6120
AGIGIATAGCATATGCGGAGACCCATGATCAGgtattttaaatttatttctacaactaaa	6180
taatteteagaagaagaagaagaagaagaagaagaagaagaagaaga	
taattctcagaacaattgttagatagaatccaaatatatacgtcctgaaagtataaaagt	6240
acttattttcgccatgggccttcagaatattggtagccgctgaatatcatgataagttat	6300
ttatccagtgacatttttatgttcactcctattatgtctgctggatacagTCTATTGTTG	6360
GTGACAAGACCATTTCCATTTTCTCCTTAATTCCTTCCTTAATTCCTTCCTTAATTCCTTCCTTAATTCCTTCCTTAATTCCTTCCTTAATTCCTTCCTTAATTCCTTCCTTAATTCCTTCCTTAATTCCTTCTCTCTTCT	
GTGACAAGACCATTGCATTTCTCCTAATGGACAAAGAGATGTATTCTGGCATGTCTTGCT D K T I A F L L M D K E M Y S G M S C L	6420
TOACAGATGCTTCTCCTGTTGATCGAGGAATTCCCCTTCACAACCEE	6400
	6480
tattgcattttaaggttcatataggttagccacggaaaatctcactctttgtgaggtaac	6540
cagggttctgatggattattcaattttctcgtttatcatttgtttattcttttcatgcat	6600
tgtgtttctttttcaatatccctcttatttggaggtaatttttctcatctattcactttt	6660
agettetaaccacachtchtechteraches comments comments	
agcttctaaccacagATGATCCATTTTTTCACAATGGCCTTGGGAGGGGGGGGGG	6720
AATTICATGGGTAACGAGgtatgtcttacatctttagatattttgtgataattacaatta N F M G N E	6780
gtttggcttacttgaacaagattcattcctcaaaatgacctgaactgttgaacatcaaag	6840
gggttgaaacatagaggaaaacaacatgatgaatgtttccattgtctagggatttctatt	6900
atgttgctgagaacaaatgtcatcttaaaaaaaacattgtttacttttttgtagtataga	6960
agattactgtatagagtttgcaagtgtgtctgttttggagtaattgtgaaatgtttgatg	
	7020
aacttgtacagTTTGGCCATCCTGAGTGGATTGACTTCCCTAGAGAGGGCAATAATTGGA F G H P E W I D F P R E G N N W S	7080
GTTATGACAAATGTAGACGCCAGTGGAACCTCGCGGATAGCGAACACTTGAGATACAAGG	7140
ttcaagtattttgaatcgcagcttgttaaataatctagtaatttttagattgcttacttg	7200
gaagtctacttggttctggggatgatagctcatttcatcttgttctacttattttccaac	
	7260
cgaatttctgattttgtttcgagatccaagtattagattcatttacacttattaccgcc	7320
tcatttctaccactaaggccttgatgagcagcttaagttgattctttgaagctatagttt	7380
caggctaccaatccacagcctgctatatttgttggatacttaccttttctttacaatgaa	7440
gtgatactaattgaaatggtctaaatctgatatctatatttctccgtctttcctcccct	7500

	• •								_				
12345678	10 90123	45678	20 9012	34567	30 78901	2345	678	40 9012	3456	50 2890	1234	60	2
						- Y - Y	0.0	7012	7770	7030	14,74.	10/09	
catgatga		F.	M N	A	FI	אר	Δ	M	NI C		D 1	7 12	
بليس لا كالمكابلية		Camos	0011					1.7	14 2	יי	נ ע	EK	
TTCTCATT F S F	L A	CATCA S	GGAA G K	AACAC O)ATA£ 1 I	TAAC 'S	CAG	CATG M	GATG	ATGA	TAAT	AAGgta	7620
aaatratr	+===	++~					_	**	ט ט	ע	14 1	Λ.	
aaatcatc	Laday	LLyad	agtg	ccggg	JEEE	itgaa	gtg	cttt	aatt	ctat	ccaa	gacaa	7680
gragaac	CHHHH	tacct	+			. 							
gtagaaac													
ctggtcaa	attco	gtaat	aget	atact	ratt	27+	30t						
ctggtcaa										3.7	37	37 E	
TGAACGTG	GTGAC	CTGGT	'ATTT	GTATI	CAAC	ككلك	אררו	מממר	CAAC	ית מיין מי	3003		7060
EKG	ע	ע ע	F '	V F	N	F H	Þ	ĸ	N	ጥ ህ	-	C	
tatatgtt	ttact	tatcc	atga	aatta	ttac	teta	ctt	~++	++==	+~+~	a= ~		- 7000
tatatgtt													
tttatgga	gaagt	aactg	aaac	aaatc	attt	tcac	atto	otct	aatt	taac	+~++		7980
•													
gatecteg	carga	Cyaaa	acag	GTATA	laagi	TGGA	TGT	GACT	TGCC	AGGG	AAGT)	CAGAC	8040
				YK	v	G	C 1	D I	. p	G	v v	10 1	,
TTGCACTG		10010	C110	GWWI	1160	TGGC	CAT	ΑΑن	GAgt	aagg	attt	gcttga	a 8100
A 1,	ם ט	D A	. W	E F	`G	G	H (G R					
ataacttt	-gaca	acaag	acaa	cayac	gray	ggca	cag	CECE	CCCa	ccaa	aaaga	actg	8160
	_												
aattgtct	catcc	atctt	tagt	tgtat	aaga	tato	cga	ctgt	ctga	gttc	ggaag	jtgtti	8220
gagcctcc	tgccc	tcccc	ctgc	gttgt	ttag	ctaa	ttc	aaaa	agga	gaaa	actg	ttatt	8280
~~+~~+						•							
gatgatct										ጥ . ር	***	D W	
TGACCATT	יבטבטיד	ጥሮልሮሮ	א מים מ	ייי או אוריי	13 CCIT	-		~~~				· · · · ·	
TGACCATT D H F	1.	SP	E (G I	P	G V	, Þ	E	T	NE	NT	G D	
TCCAAATT	CCTTC	AAAGT	CCTC	ייייי	TCC	מ מסים	ርእጥ	ساست					- 0466
PNS	F.	K V	ъ :	5 P	A	R T	`С	v			-		•
tgacctcc	ctttt	tattg	tggti	tttgt	tcat	agtt	atti	tgaa	taca	ataq	aagti	aact:	a 8520
ttgattac	cgcca	caatc	gcca	gttaa	gtcc	tctg	aact	tact	aatt	tgaa	aggta	aggaat	8580
agccgtaa	taagg	tctac	tttt	racat	ctta	~+ ~+	+						
-99	99			ggcac	ccca	cege	Lac	aaaa	caaa	agga	tgcc	aaaaa	a 8640
attettete	ctatc	ctctt	tttc	cctaa	acca	gtgc	atgi	tagc	ttgc	acct	gcata	aacti	8700
aggtaaat	gatca	aaaat	gaagi	ttgat	ggga	actt	aaaa	accg	ccct	gaag	taaag	gctag	8760
aatagtca	tataa	tgtcc	accti	ttggt	gtct	gcgc	taad	catc	aaca	acaa	cata	cctcgt	8820
	_												
gtagtccc	acaaa	grggt	ttca	39999	aggg	taga	gtgl	tatg	caaa	actt	actc	ctatc	8880
	~~~~	~=+-		k									
cagaggtag	yayag	yaitt	LEEC	aacag	accc	rtgg	CEC	aaga	aaaa	aagt	ccaa	aaaga	a 8940
ntaacene:	antes	22000											
gtaacagaa	-yrya	aayca	مدمدز	gryra	ycta	.aagc	gac	ccaa	CEEG	ttta	ggact	tgaag	9000

	10 20 30 40 50 60 34567890123456789012345678901234567890	
9060	ttgttgttgttgaaacagtgcatgtagatgaacacatgtcagaaaatggacaacacag	
	attttgtgcaagtcaaaaaaatgtactactatttctttgtgcagctttatgtatagaa	
	gttaaataactaatgaattttgctagcagaaaaatagcttggagagaaatttttata	
9240	gaactaagctaactatattcatctttctttttgcttcttcttctccttgtttgt	
9300	TTATTACAGAGTTGATGAACGCATGTCAGAAACTGAAGATTACCAGACAGA	
03.60	TGAGCTACTACCAACAGCCAATATCGAGGAGAGTGACGAGAAACTTAAAGATTCGTTA	
9360	E L L P T A N I E E S D E K I K D c i	
9420	TACAAATATCAGTAACATTGACGAACGCATGTCAGAAACTGAAGTTTACCAGACAGA	
	T N I S N I D E R M S E T E V Y Q T D	
9480	TTCTAGTGAGCTACTACCAACAGCCAATATTGAGGAGAGTGACGAGAAACTTAAAGAT SSELLPTANIESSDEKIKD	
0540	GTTATCTACAAATATCAGTAACATTGATCAGACTGTTGTAGTTTCTGTTGAGGAGAGA	
	L S T N I S N I D O T V V V S V F F B	
9600	CAAGGAACTTAAAGATTCACCGTCTGTAAGCATCATTAGTGATGTTGTTCCAGCTGAA	
	K E L K D S P S V S I I S D V V P A E	
9660	GGATGATTCAGATGCAAACGTCTGGGGTGAGGACTAGTCAGATGATTGAT	
9720	ACCGATTGGTGATCGCTATCCTTGCTCTCTGAGAAATAGGTGAGGCGAAACAAAAAAT	
9780	TTTGCATGATAAAAAGTCTGATTTTATGATCGCTATCCTCGCTCTCTGAGAAAGAA	
9840	AACAAAGGCGACTCCTGGACTCGAATCTATAAGATAACAAAGGCGACTCCTGGGACTC	
9900	ATCTATAAGATAACAAAGGCAATTCCAAGACTTGAATCTATAAAAATTTAGTTAAGA	
9960	GATTAACGTCCGATCCTAATTCGAATCGAGGCATCTTACCACTCCATTGATAATTATA	
•	ACTICA ATTA ACTICA TIA TIA ACTICA TIA TIA ACTICA AC	
10020	AGTCAATAAGTCATATAAWAGTATTAAAAACTAAATTGACTTGATCGGTCTATCAAAA	
10080	AGATMAAATTGTGTTCATATGTAACATTTTTGTTGTCACAATTAGCTTAATTACATC	
10140	CATGTGCAATAACAAAGAAATGATAGGAATTTAGAGATTCCAATTTTTTTGTTGCCA	
10200	ATTAACTTAATTACATCTTTCATTTGCAATAACAAAGAAATGATAGGAATTTAGAGAT	
10260	AGTGTCAATACACAACCTAGGCCAACATCGAAAGCATAACTGTAAACTCATGCATG	
10320	ATCAGTCGTAAAAATGAATAAATGCGACATAAAAACAAATTGCATGTATCATTAATG	
10380	ACTTAACTACAAGTAAAAATAAATTTAACAAATGTAACTTAACTACAAGTAAAAATAA	
10440	PGCTTCTATCATTAACAAACAAACAGAATTAAAAAGAAAAAAAA	
10500	CATTCGATAAAAAAAAATACCAAATTCATAATGCAAGGAAAACGAAACGCGTCCTGA	,

<del></del>						•	
1234567	10 8901234	20 5678901234	30 5678901234	40 5678901234	50 56789012345	60 67890	
TCGGGTA	TCAACGA	TGAAATGGAC	CAGTTGGATC	GACTGCCTGC	ACAACGTTAGG	TATGC	10560
САААААА	AAGAACA	CGATCCTTTG	CACCCGTTCG	ATGATTATCA(	GTATGTTCACA	AAAAA	10620
AACTTAA	GTTCATC	CCAGTGTACA	ACAGCCCCAA	CATCTGCCCC	AAGTAACAAAA	AACAA	10680
CCAATTT	ATCTTAT	TCTTATCTGC	CACAAAATAA	TCGGTTTCAC	ACTATTCTCTT	GTTAT	10740
ACAAAAT	TGACAAG	TAGGAAGGAG	AGGAGTCATC	CAAATAAACG	STGCACGTTCT	ITGAG	10800
AAAAGTC	TTTTTTT	TCGTAAGATC	CAATTTCAAC	AAACTTTTCT	CAAGTCAAAA'	TTCCT	10860
GATAGTG!	TATCTCC	TCTCGACGAC	CTCTTGCATT	GAACGATCTC	CGCTTATCATG	AAAAG	10920
TTGCTTG	GATAACA	AGTATTGCAA	GGGGGGACA	GTAGCTATTA	AGTTAGTCGGC	CCAAG	10980
GAAATGG	AGGAGTG	ATAGTCTCGA	ATATTATTCA	CCTCTTTAGC	ATTACCCGGTC	IGGCT	11040
TTAAGGAG	STTACGT	CTTTTACGCT	CGCCAATTTC	TTTTTTTAGAZ	ATGGTTGGTGT(	CAAAA	11100
TCGCGAGT	TGTGGA	AGGTTCAAGT	PACTCGATTC	GTGATTTTCA	AGTATGAGTGG	rgaga	11160
GAGATTC	TTTATAE	TCACGAGGTGT	TATTCGAGGT	CTAGTAGAACC	SAAGGGTGTCA	TAAT	11220
GAAAGTTT	CAAGAG	TTCATCATCAT	CTTCTTCTA	GTAGATTTTC	CTTTCAAATG	AGTAT	11280
GAAAATTO	CTTCCTC	TTTTCTATTG	ATTTTCTTCA	TTGTTTTCTTC	ATTGTTGTGG	PTGTT	11340
ATTGAAAA	GAAAGA	AAATTTATAAC	`AGAAAAAGA'	TGTCAAAAAA	AGGTAAAATG	AAAGA	11400
GTATCATA	TACTTA	AAGAGTTGCGT	ragagataag	TCAAAAGAAAC	AGAATTATAG:	TAATT	11460
TCAGCTAA	GTTAGA	ATTC					11478

. 1.5₹




Fig 9




Fig 10

# PLI/EP96/03053

## INDICATIONS RELATING TO A DEPOSITED MICROORQANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism	referred to in the description				
on page 19 , lines 1					
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet				
Name of depositary institution					
The National Collections of Industria	l and Marine Bacteria Limited (NCIMB)				
Address of depositary institution (including postal code and cour					
23 St. Machar Drive					
Aberdeen					
Scotland AB2 1RY					
United Kingdom					
Date of deposit	Accession Number				
13 July 1995					
C. ADDITIONAL INDICATIONS (leave blank if not applied	NCIMB 40754, NCIMB 40751, NCIMB 40752				
C. ADDITIONAL INDICATIONS (leave blank if not applied	cable) This information is continued on an additional sheet				
grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample. (Rule 28(4) EPC).					
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)					
·					
E SEPARATE EURNISHING OF INDICATIONS					
E. SEPARATE FURNISHING OF INDICATIONS (le					
Number of Deposit")	al Bureau later (specify the general nature of the indications e.g., "Accession				
. <i>'</i>					
For analysis of Control					
For receiving Office use only	For International Bureau use only				
This sheet was received with the international application	This sheet was received by the International Bureau on:				
Authorized officer					
P.L.R. Petter	Authorized officer				
<u> </u>					
rm PCT/RO/134 (July 1992)	. I <u> </u>				

## BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF THE DEPOSIT OF MICROCRGANISM FOR THE PURPOSES OF PATENT PROCEDURE

PCT/EP 96 / 03 05 3

Danisco Biotechnology : gebrogade 1 P O Box 17 DK-1001 Copenhagen K Denmark INTERNATIONAL FORM

RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT issued pursuant to Rule 7.1 by the INTERNATIONAL DEPOSITARY AUTHORITY identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISH	
Identification reference given by the DEPOSITOR:	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY:
Escherichia coli DH5∢-pBEAll	NCIMB 40754
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED	TAXONOMIC DESIGNATION
The microorganism identified under I above	was accompanied by:
a scientific description	
X a proposed taxonomic designation	
(Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority according which was received by it onl3 July 1995 (c	cepts the microorganism identified under I above, late of the original deposit)
IV. RECEIPT OF REQUEST FOR CONVERSION	
The microorganism identified under I above Depositary Authority on (day a request to convert the original deposit (was received by it on	te of the original deposit) and
V. INTERNATIONAL DEPOSITARY AUTHORITY	
Name: NCIMB Ltd  23 St Machar Drive Aberdeen Scotland Address: UK AB2 1RY	Signature(s) of person(s) having the power to represent the International Depositary Authority or of authorized official(s):  Date: 21 July 1995 Teence Dando

Form BP/4 (sole page)

where Rule 6.4(d) applies, such date is the date on which the status of international depositary authority was acquired.

# BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF THE DEPOSIT OF MICROORGANISMS FOR THE PURPOSES OF PATENT PROCEDURE

Danisco Biotechnology Langebrogade 1 P O Box 17 DK-1001 Copenhagen K Denmark PCT/EP96/03053

INTERNATIONAL FORM

VIABILITY STATEMENT issued pursuant to Rule 10.2 by the INTERNATIONAL DEPOSITARY AUTHORITY identified on the following page

NAME AND ADDRESS OF THE PARTY TO WHOM THE VIABILITY STATEMENT IS ISSUED

I. DEPOSITOR	II. IDENTIFICATION OF THE MICROORGANISM
Name: As above	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY:
Address:	NCIMB 40754 Date of the deposit or of the transfer:
	13 July 1995
III. VIABILITY STATEMENT	
The viability of the microorganism	identified under II above was tested
on 13 July 1995	² . On that date, the said microorganism was
X viable	
no longer viable	

Indicate the date of the original deposit or, where a new deposit or a transfer has been made, the most recent relevant date (date of the new deposit or date of the transfer).

In the cases referred to in Rule 10.2(a)(ii) and (iii), refer to the most recent viability test.

Mark with a cross the applicable box.

/. CONDITIONS UNDER WHICH THE VIABILITY TEST H.	AS BEEN PERFORMED
V. INTERNATIONAL DEPOSITARY AUTHORITY	
Name:  NCIMB Ltd  Address: 23 St Machar Drive Aberdeen Scotland 11K AR2 1RY	Signature(s) of person(s) having the power to represent the International Depositary Authority or of authorized official(s):  Terence Dando  Date:  21 July 1995

Fill in if the information has been requested and if the results of the test were negative.

## BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF THE DEPOSIT OF MICROORGANISMS FOR THE PURPOSES OF PATENT PROCEDURE

DR A Buchter-Larsen
Panisco Biotechnology
Ingebrogade 1
P O Box 17
DK-1001 Copenhagen

Denmark

INTERNATIONAL FORM

PCT/EP 96 / 03 05 3

RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT issued pursuant to Rule 7.1 by the INTERNATIONAL DEPOSITARY AUTHORITY identified at the bottom of this pag.

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the Accession nu DEPOSITOR: INTERNATION	umber given by the NL DEPOSITARY AUTHORITY:
Bacteriophage $\lambda$ EMBL3 SP6/T7 $\lambda$ SBE3.2 NCIM	B 40751
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DES	SIGNATION
The microorganism identified under I above was accompanie	ed by:
a scientific description	
X a proposed taxonomic designation	
(Hark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	·
This International Depositary Authority accepts the micr which was received by it on 13 July 1995(date of the or	oorganism identified under 1 above, iginal deposit) 1
IV. RECEIPT OF REQUEST FOR CONVERSION	
beposit to a deposit to	(dinal debosic) and
V. INTERNATIONAL DEPOSITARY AUTHORITY	
to represe	s) of person(s) having the power nt the International Depositary or of authorized official(s):

Form BP/4 (sole page)

Where Rule 6.4(d) applies, such date is the date on which the status of international depositary authority was acquired.

#### BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF THE DEPOSIT OF MICROORGANISMS FOR THE PURPOSES OF PATENT PROCEDURE

Dr A Buchter-Larsen
Danisco Biotechnology
Langebrogade 1
P O Box 17
DK-1001 Copenhagen K
Denmark

INTERNATIONAL FORM

VIABILITY STATEMENT issued pursuant to Rule 10.2 by the INTERNATIONAL DEPOSITARY AUTHORITY identified on the following page

NAME AND ADDRESS OF THE PARTY TO WHOM THE VIABILITY STATEMENT IS ISSUED

I. DEPOSITOR	II. IDENTIFICATION OF THE MICROORCANISM
Name: As above Address:	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: NCIMB 40751 Date of the deposit or of the transfer:
	13 July 1995
III. VIABILITY STATEMENT	
The viability of the microorganism identified un on 19 July 1995	der II above was tested  2. On that date, the said microorganism was
X viable  no longer viable	

Indicate the date of the original deposit or, where a new deposit or a transfer has been made, the most recent relevant date (date of the new deposit or date of the transfer).

In the cases referred to in Rule 10.2(a)(ii) and (iii), refer to the most recent viability test.

Mark with a cross the applicable box.

tv.	CONDITIONS UNDER WHICH THE VIABILITY TEST	HAS BEEN PERFORMED
ļ .		
		· · · · · · · · · · · · · · · · · · ·
v.	INTERNATIONAL DEPOSITARY AUTHORITY	
Nam Add	Press: 23 St Machar Drive Aberdeen Scotland LN AB2 1RY	Signature(s) of person(s) having the power to represent the International Depositary Authority or of authorized official(s):  Date: 20 July 1995

⁴ Fill in if the information has been requested and if the results of the test were negative.

# BUDAPEST TREATY CH THE INTERNATIONAL RECOGNITION OF THE DEPOSIT OF MICROORGANISMS FOR THE PURPOSES OF PATENT PROCEDUP.

Dr A Buchter-Larsen
Danisco Biotechnology
Langebrogade 1

, Box 17
DK-1001 Copenhagen
Denmark

INTERNATIONAL FORH

PCT/EP 96 / 03 05 3

RECEIPT IN THE CASE OF AN ORIGINAL DEPOSITIONS of the INTERNATIONAL DEPOSITARY AUTHORITY identified at the bottom of this page

NAME AND ADDRESS OF DEPOSITOR

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR:	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY:
Bacteriophage & EMBL3 SP6/T7 &SBE3	3.4 NCIMB 40752
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSE	ED TAXONOMIC DESIGNATION
The microorganism identified under I above	e was accompanied by:
a scientific description	
A proposed taxonomic designation	
(Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority a which was received by it on13 July 1995	occepts the microorganism identified under I above, sidate of the original deposit)
IV. RECEIPT OF REQUEST FOR CONVERSION	
The microorganism identified under I about Depositary Authority on a request to convert the original deposit was received by it on	idate of the oficinal ceposit, and
V. INTERNATIONAL DEPOSITARY AUTHORITY	
Name: NCIMB Ltd 23 St Machar Drive Aberdeen Scotland	Signature(s) of person(s) having the power to represent the International Depositary Authority or of nutberized official(s):
Aberdeen Scottano	Date: 20 July 1995

form BP/4 (sole page)

Where Rule 6.4(d) applies, such date is the date on which the status of international depositary authority was acquired.

#### PCT/EP96/03053

BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF THE DEPOSIT OF MICROORGANISMS FOR THE PURPOSES OF PATENT PROCEDURE

Dr A Buchter-Larsen
Danisco Biotechnology
Langebrogade 1
P O Box 17
DK-1001 Copenhagen K
Denmark

INTERNATIONAL FORM

VIABILITY STATEMENT issued pursuant to Rule 10.2 by the INTERNATIONAL DEPOSITARY AUTHORITY identified on the following page

NAME AND ADDRESS OF THE PARTY TO WHOM THE VIABILITY STATEMENT IS ISSUED

I. DEPOSITOR	II. IDENTIFICATION OF THE MICROORGANISM			
Name: As above Address:	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: NCIMB 40752 Date of the deposit or of the transfer:			
	13 July 1995			
III. VIABILITY STATEMENT				
The viability of the microorganism identified under II above was tested on 19 July 1995 2. On that date, the said microorganism was				
X viable				
no longer viable				

Indicate the date of the original deposit or, where a new deposit or a transfer has been made, the most recent relevant date (date of the new deposit or date of the transfer).

In the cases referred to in Rule 10.2(a)(ii) and (iii), refer to the most recent viability test.

Mark with a cross the applicable box.

tv.	CONDITIONS	UNDER WHICH	THE VIABILITY	TEST H	AS BEEN	PERFORMED *
		•				•
			•			
						-
					<u> </u>	
v.	INTERNATIO	NAL DEPOSITA	RY AUTHORITY			
Mam Add	ress: 2	NCIMB   3 St Machar perdeen So	Drive cotland		Sig to Aut Dat	nature(s) of person(s) having the power represent the International Depositary hority or of authorized official(s):  e:  20 July 1995

⁴ Fill in if the information has been requested and if the results of the test were negative.